

TEXAS MEDICATION ALGORITHM PROJECT PROCEDURAL MANUAL

SCHIZOPHRENIA TREATMENT ALGORITHMS

Tami R. Argo, PharmD, MS, BCPP


M. Lynn Crismon, PharmD

Alexander L. Miller, MD

Troy A. Moore, PharmD, MS, BCPP

Sherrie D. Bendele, BS

Brandon Suehs, PharmD



Schizophrenia Algorithm Procedural Manual

Table of Contents

Disclaimer	1
Author Affiliations	2
Financial Disclosures	3
Overview of the Texas Medication Algorithm Project	4
Clinical Management	6
At-a-Glance Schizophrenia Medication Algorithm	7
Schizophrenia Algorithm	8
Description of Algorithm Stages	11
Tactics and Critical Decision Points	13
Process Measures: Evaluation of Patient Response	15
Positive Symptom Rating Scale (PSRS) and Brief Negative Symptom Assessment (GNSA)	17
Clinician Ratings.....	17
Patient Self-Ratings	18
Medications and Dosing	19
Medication Discontinuation and Maintenance	20
Medication Discontinuation.....	20
Medication Maintenance.....	20
Documentation	21
Outpatient Documentation.....	21
Inpatient Documentation.....	22
Modifications for Inpatient Use	23
Inpatient to Outpatient Transition	25
Outpatient to Inpatient Transition	26

Appendix A: Process Measures	27
4-ITEM POSITIVE SYMPTOM RATING SCALE (PSRS).....	28
BRIEF NEGATIVE SYMPTOM ASSESSMENT SCALE (version 5.0)	32
SCORE SHEET for 4-ITEM POSITIVE SYMPTOM RATING SCALE AND BRIEF NEGATIVE SYMPTOM ASSESSMENT .	35
Appendix B: Communications	38
Appendix C: Medication Charts	39
Antipsychotics, Second Generation	40
Antipsychotics, First Generation.....	42
Adjunctive Agents, Agitation/Insomnia.....	44
Adjunctive Agents, Insomnia.....	44
Adjunctive Agents, Aggression/Hostility	45
Adjunctive Agents, Depression.....	47
ADDITIONAL REFERENCES FOR DRUG INFORMATION.....	51
Appendix D: Side Effect and Co-Existing Symptom Management	52
Management of Associated or Co-Existing Symptoms in Schizophrenia.....	52
Management of Treatment-Emergent Side Effects in Schizophrenia	54
Appendix E: TMAP Publications.....	57
Appendix F: Minimum Data Set for Documentation	60

Disclaimer

This manual is based upon the evidence based, expert consensus recommendations presented in the article:

Moore TA, Buchanan RW, Buckley PF, Chiles JA, Conley RR, Crismon ML, Essock SM, Finnerty M, Marder SR, Miller DD, McEvoy JP, Robinson D, Schooler NR, Shon SP, Stroup TS, Miller AL. The Texas Medication Algorithm Project Antipsychotic Algorithm for Schizophrenia: 2006 Update. *J Clin Psychiatry*. 2007; 68:1751-1762.

The manual also reflects the experiences of the TMAP team in conducting the research evaluating use of the algorithms, as well as in implementing the algorithms in public mental health systems. These algorithms reflect the state of knowledge, current at the time of publication, on effective and appropriate care as well as clinical consensus judgments when research-based knowledge is lacking. The inevitable changes in the state of scientific information and technology mandate that periodic review, updating, and revisions will be needed. These guidelines (algorithms) may not apply to all patients, and each must be adapted and tailored to each individual patient. The authors bear no responsibility for the use and/or modification of these guidelines by third parties. The provision of clinical care, including recommendations contained in these or other guidelines, in whole or in part, is entirely the responsibility of the clinician.

The Texas Medication Algorithm Project (TMAP) schizophrenia algorithm and this manual are copyrighted by the Texas Department of State Health Services (DSHS). If you are using or adapting the entire manual, sections, tables or figures, please contact us for written permission. Contact information can be found in Appendix B. Please use proper citation and acknowledgement of the authors and this manual when citing or referencing the manual:

Argo TR, Crismon ML, Miller AL, Moore TA, Bendele SD, Suehs B. *Texas Medication Algorithm Project Procedural Manual: Schizophrenia Algorithm*. The Texas Department of State Health Services. 2007.

Author Affiliations

M. Lynn Crismon, PharmD, BCPP

Dean

James T. Doluisio Chair and Behrens Inc. Centennial Professor

College of Pharmacy

The University of Texas at Austin

Austin, TX

Tami R. Argo, PharmD, MS, BCPP

Clinical Assistant Professor

College of Pharmacy

The University of Texas at Austin

Austin, TX

Alexander L. Miller, M.D.

Professor

Department of Psychiatry

The University of Texas Health Science Center at San Antonio

San Antonio, TX

Troy A. Moore, PharmD, MS, BCPP

Assistant Professor

Department of Psychiatry

The University of Texas Health Science Center at San Antonio

San Antonio, TX

Sherrie D Bendele, BS

Program Coordinator

College of Pharmacy

The University of Texas at Austin

Austin, TX

Brandon T Suehs, PharmD, PhD candidate

Graduate Research Assistant

College of Pharmacy

The University of Texas at Austin

Austin, TX

Financial Disclosures

Dr. Crismon has received grant/research support from Eli Lilly, Forest, Janssen and Shire; and is on the speakers/advisory board of Corcept Therapeutics, Cyberonics, Eli Lilly, Forest, Janssen, and Shire.

Dr. Argo has no significant financial relationships to disclose.

Dr. Miller has received grant/research support from Alexza, AstraZeneca, Bristol Myers Squibb, Eli Lilly, Forest Pharmaceuticals, Glaxo Smith Kline, Janssen Research Foundation, Organon, and Pfizer; has served as a consultant to AstraZeneca, Bristol Myers Squibb, Centene, Dainippon Sumitomo, Eli Lilly, Janssen Research Foundation, Organon, Pfizer, and Solvay.

Dr. Moore has no significant financial relationships to disclose.

Ms. Bendele has no significant financial relationships to disclose.

Dr. Suehs has no significant financial relationships to disclose.

Overview of the Texas Medication Algorithm Project

Algorithms facilitate clinical decision making by providing clinicians with large amounts of current information on the newest psychotropic medications and research data, as well as specific treatment sequences with tactical recommendations. Patients receive the benefit of patient education, which should enhance adherence to the treatment program. Algorithms are designed with the objectives of long-term safety, tolerability, and full symptom remission — not just response. The employment of such treatment guidelines to assertively treat the severely and persistently mentally ill (SPMI) population may enhance patient outcomes while improving the utilization of crisis/hospital services and improving accountability for scarce resources — thereby increasing the overall efficiency of patient care.

Beginning in 1995, The Texas Medication Algorithm Project (TMAP) was developed by the Texas Department of Mental Health and Mental Retardation (TDMHMR¹) in collaboration with Texas universities to assess the value of an algorithm-driven disease management program in the pharmacological management of mentally ill patients. The result has been a set of algorithms for the treatment of the three major disorders most commonly encountered in the Texas public mental health system: schizophrenia (SCZ), bipolar I disorder (BDI), and major depressive disorder (MDD). A best practice treatment has been defined as a series of treatment steps that guides physicians in determining medication treatment plans, thereby generating the best outcome for each individual consumer. The algorithms consist of both treatment strategies (recommended sequential medication regimen options) and treatment tactics (recommended options for optimal use of a medication regimen in a given patient). Equal attention should be given to the treatment tactics as to the strategies. A critical component of all the algorithms in TMAP is measurement and documentation of target symptoms with quantitative scales.

Practitioners, patients, families, and administrators all contributed to the formulation and implementation of TMAP, ensuring an optimum level of efficacy and practicality. Phase 1 of TMAP dealt with the development of these algorithms using expert consensus. In Phase 2, the feasibility of algorithm implementation in the TDMHMR system was evaluated. Phase 3 evaluated the clinical and economic impact of medication treatment algorithms for MDD, SCZ, and BDI in comparison with Treatment As Usual (TAU). Please refer to Appendix F for a list of publications.

Implementation of the algorithms on a system wide basis was the next step in offering high quality care to the SPMI patient population in the public mental health sector. This rollout was referred to as Texas Implementation of Medication Algorithms (TIMA) (Phase 4 of TMAP) in order to distinguish it from the research phases of TMAP. However, in order to retain name identity, TMAP is once again being used for the program. The rollout began with the training of physicians and support personnel in algorithm implementation.

Continued revision may be required in the structure and function of clinical staff to increase patient education and adherence, to improve follow up, and to develop psychosocial supports to improve symptom recognition, symptom control, and functional restoration. Continuous education, consultation, and collaboration are necessary for both clinicians and administrators in making timely revisions in clinical procedures and budgetary allocations. From a clinical and administrative perspective, medication algorithms should demonstrate validity with far-reaching and long-term applications.

¹ State public mental health services are now provided as a component of the Texas Department of State Health Services (DSHS).

For additional information regarding the development of the most current Schizophrenia Algorithm, please refer to the article: Moore TA, Buchanan RW, Buckley PF, Chiles JA, Conley RR, Crismon ML, Essock SM, Finnerty M, Marder SR, Miller DD, McEvoy JP, Robinson D, Schooler NR, Shon SP, Stroup TS, Miller AL. The Texas Medication Algorithm Project Antipsychotic Algorithm for Schizophrenia: 2006 Update. *J Clin Psychiatry*. 2007; 68:1751-1762.

Clinical Management

- At baseline and throughout treatment, the patient should be evaluated for possible psychosocial interventions. Psychosocial rehabilitation programs oriented toward improving patients' adaptive functioning are the mainstay of nondrug treatment for schizophrenia. These programs may include basic living skills, social skills training, basic education, work programs, and supported housing.
- Use of the algorithms assumes that the clinician has made a thorough evaluation and an accurate diagnosis. If a patient completes trials of two stages of the algorithm without observable positive outcomes, the patient should be re-evaluated for accuracy of diagnosis and the occurrence of co-occurring general medical and mental disorders, including substance abuse.
- If co-occurring substance abuse is present, concomitant treatment of both the schizophrenia and the substance abuse disorder must be implemented in order to obtain optimal positive patient outcomes.
- Brief symptom ratings (PSRS, BNSA) should be completed at each medication visit so that medication treatment decisions are guided by objective data.
- Adequate documentation should be completed for each algorithm stage and treatment choice (i.e., critical decision points). If algorithm stages are skipped or if treatment is different from the algorithms, the rationale should be adequately documented.
- The frequency of clinic visits should be adequate to monitor for symptom changes and adverse effects, to adjust doses as necessary to achieve an optimum therapeutic trial, and change regimens when suboptimal clinical response is observed after regimen optimization.
- When a choice exists between brand, generic, or different formulations (e.g., slow release) of a recommended medication, always initiate treatment with the form that is likely to be best tolerated by the patient, which will lead to enhanced adherence with treatment. Careful attention should be given to adequate dose and duration of treatment for each chosen regimen.
- All patients with schizophrenia who achieve a satisfactory clinical response (minimal positive symptoms and preferably remission) should continue treatment indefinitely as maintenance treatment.
- If medication acquisition cost is a consideration in medication selection, these decisions should be addressed within a specific treatment stage. If all other things are equal (i.e., efficacy, safety, tolerability), then a less expensive medication regimen within a specific algorithm stage may be considered.

At-a-Glance Schizophrenia Medication Algorithm

At-a-Glance Schizophrenia Medication Algorithm

Visit Frequency: While medications are being actively adjusted, patients should be seen every 2 weeks. As medications are stabilized and patients exhibit stable, positive response, visit intervals can be gradually lengthened to every 4 weeks. When patients achieve a stable response, visit frequency can be scheduled for every 8-12 weeks, as individually determined. Additional patient contact (e.g., by telephone) may be necessary to provide optimal care for a symptomatic patient.

Assessment Frequency: The 4-Item Positive Symptoms Rating Scale (PSRS) and the Brief Negative Symptom Assessment (BNSA) should be completed at each medication visit.

Criteria for Medication Change: Medication changes are made after evaluation of tolerability, efficacy across multiple symptom domains, and safety. Clinicians should consult the Tactics and Critical Decision Points for the Treatment of Schizophrenia after review of symptom patterns and severity on the Process Measures Graph, as well as any medication side effects and tolerability. The goals of treatment are symptomatic remission (or minimal positive symptoms), return of psychosocial functioning, and prevention of relapse and recurrence.

As much as possible, patients should receive an adequate trial of each antipsychotic. Patients need at least four weeks of therapeutic doses of an antipsychotic (excluding clozapine) before they can be classified as “non-responders” to the medication. Clozapine requires more time, up to three months. Assessing the full effects of an antipsychotic can take 12 weeks or longer. During acute relapses, multi-week trials of agents are difficult to sustain. However, failure to respond to an antipsychotic in 1-2 weeks should not eliminate it from future consideration as a possibly effective agent. Another trial may be worthwhile under more symptomatically stable circumstances.

Evaluations: At each medication visit, a physician will assess core symptom severity, overall functional impairment, and side effect severity. The provider/physician can complete the PSRS, BNSA and have the patient complete the global self-rating of symptom severity and side effects.

Medication Doses: Appropriate dosage ranges for medications used in the algorithms are included in *Appendix C*. Doses outside of the ranges should have a chart note indicating “change from algorithm recommended” and documentation of rationale for change. Doses above the usual therapeutic range should be time limited (e.g., 4-6 weeks), and response to this dose evaluated using the brief clinical ratings. If improvement has not occurred with the higher than usual dosage in this time frame, then treatment should be changed to the next treatment stage.

Documentation: Uniform documentation is an important component of the algorithm program. Clinical rating scale information, response to treatment, prescribed medications, and the rationale for changing medications should be clearly documented on the Clinical Report Form.

Schizophrenia Algorithm

Choice of antipsychotic (AP) should be guided by considering the clinical characteristics of the patient and the efficacy and side effect profiles of the medication.
Forward stage(s) can be skipped depending on clinical picture or history of antipsychotic failures, and returning to an earlier stage may be justified by history of past response.

* First episode patients usually require lower antipsychotic dosing and should be closely monitored due to greater sensitivity to medication side effects. Lack of consensus on inclusion of FGAs as option for first episode.

FGA = First generation Antipsychotic
 SGA = Second generation Antipsychotic
 ECT = Electroconvulsive Therapy

Stage 1: First Episode Schizophrenia*^a

Trial of a single SGA
 (ARIPRAZOLE, OLANZAPINE,
 QUETIAPINE, RISPERIDONE, or
 ZIPRASIDONE)

Partial or Non-response

Stage 2

Trial of a single SGA or FGA
 (not SGA tried in Stage 1)

Partial or Non-response

Stage 3

CLOZAPINE

Partial or Non-response^b

Stage 4

CLOZAPINE
 +
 (FGA, SGA or ECT)

Non-response

Stage 5

Trial of a single agent
 FGA or SGA
 (not tried in Stages 1 or 2)

Stage 6

Combination Therapy
 E.g. SGA + FGA, combination of SGAs,
 (FGA or SGA) + ECT, (FGA or SGA) +
 other agent (e.g. mood stabilizer)^c

Consider earlier trial of clozapine in patients with a history of recurrent suicidality, violence, or comorbid substance abuse. Persistence of positive symptoms > 2 years warrants and > 5 years requires a clozapine trial, independent of number of preceding antipsychotic trials.

Inconsistent results in RCTs

Value in clozapine failures not established

Case reports, no controlled studies of combinations in long term treatment of schizophrenia

^a If patient is inadequately adherent at any stage, the clinician should assess contributing factors and consider a long-acting antipsychotic preparation, such as risperidone microspheres, haloperidol decanoate or fluphenazine decanoate.

^b A treatment refractory evaluation should be performed to reexamine diagnosis, substance abuse, medication adherence, and psychosocial stressors. Cognitive Behavioral Therapy and other psychosocial augmentations should be considered.

^c Whenever a second medication is added to an antipsychotic (other than clozapine) for the purpose of improving **psychotic** symptoms, the patient is considered to be in Stage 6.

Schizophrenia Algorithm

The medications algorithms were designed with the objectives of optimizing long-term safety and tolerability, and achieving full symptom remission. The recommendations contained in the algorithm are based on the evidence available at the time of their development. When evidence is not available to guide algorithm development, expert consensus drives algorithm recommendations.

First-Episode vs. Multi-Episode Schizophrenia

The schizophrenia treatment algorithm is applicable both to patients with first-episode schizophrenia and chronic, multi-episode schizophrenia. While evidence suggests that individuals with a first-episode of schizophrenia are more likely to respond to lower doses of antipsychotic and may be more susceptible to certain antipsychotic side effects, no specific recommendations regarding dosage or side effects was made by the consensus panel. The majority opinion of the consensus panel is that second generation antipsychotic medications are favored as initial treatment for first-episode schizophrenia.

Medication Change Criteria

Medication changes should only be considered after evaluation of tolerability, efficacy, and safety. As much as possible, patients should receive an adequate trial of each antipsychotic. Patients need at least four weeks of a therapeutic dosage of an antipsychotic (excluding clozapine) before they can be classified as “non-responders” to the medication. Clozapine requires more time, up to three months. Assessing the full effects of an antipsychotic can take 12 weeks or longer. During acute relapses, multi-week trials of agents are difficult to sustain. However, failure to respond to an antipsychotic in 1-2 weeks should not eliminate it from future consideration as a possibly effective agent. Another trial may be worthwhile under more symptomatically stable circumstances.

Medication Serum Concentrations:

If lithium or valproate is utilized in Stage 6, serum concentrations should be obtained in order to assist in the evaluation of medication response and tolerability. It is recommended that within 2 weeks after initiating lithium (Li) or valproate (VPA), the patient be receiving the minimum target dose. If possible, a serum concentration should be obtained approximately 5-7 days (~5 half-lives) after reaching the initial target dose (i.e. about 2-3 weeks after starting the trial). Further dosage adjustments should be made based on efficacy and tolerability, with medication serum concentration being used as needed to inform an evaluation of efficacy and tolerability. Target serum concentrations are provided in *Appendix C*. Keep in mind that these recommendations are based on data from studies in acute mania. No therapeutic blood levels have been established for use of these agents as augmentation treatments in schizophrenia.

Monitoring Atypical Antipsychotics:

Routine health monitoring is essential to detection and management of side effects that may result from treatment with antipsychotic medications. As the use of atypical antipsychotic medications has become increasingly more widespread, several health implications have been recognized through case reports, post-marketing surveillance, and pharmacoepidemiological studies. Some antipsychotic

medications have been associated with weight gain, dyslipidemia, hyperglycemia, and altered EKG findings. In addition, significant research suggests that patients with schizophrenia may be more likely to experience certain health conditions, such as diabetes, even in the absence of medication-related risk factors. Based on this information, the Texas public health system has adopted the Mount Sinai Conference health monitoring guidelines. These guidelines include recommendations for monitoring the physical health of patients receiving antipsychotic medications, including recommendations for routine monitoring of metabolic side effects such as weight gain, diabetes, and hyperlipidemia. The Mt. Sinai Conference monitoring guidelines can be accessed using the following citation:

Marder SR, Essock SM, Miller AL, et al. Physical Health Monitoring of Patients With Schizophrenia. *American Journal of Psychiatry* 2004;161:1334-49.

Co-Occurring Substance Abuse

It is common for individuals with schizophrenia to have a co-occurring substance abuse disorder. In these cases, it is extremely important that both disorders are appropriately treated in order to maximize the likelihood of a beneficial treatment outcome. The most common substances abused by patients diagnosed with schizophrenia are alcohol, marijuana and cocaine. Nicotine use is also extremely common among patients diagnosed with schizophrenia. Clozapine has demonstrated the most significant and consistent positive effect on outcomes in patients with schizophrenia and a co-occurring substance use disorder. As such, the consensus panel recommends that clozapine treatment be considered earlier in patient with a co-occurring substance use disorder.

Description of Algorithm Stages

This section of the manual explains the rationale behind the sequence of stages in the Schizophrenia Algorithm and highlights changes made at the Schizophrenia Algorithm Update Conference in June 2006.

Stage 1

Stage 1 is treatment for patients with new onset schizophrenia and includes monotherapy with a second-generation antipsychotic. Medication choices at Stage 1 include monotherapy with aripiprazole, olanzapine, quetiapine, risperidone, or ziprasidone.

Once adequate dose titration and treatment duration are achieved, decide how well the patient has responded to the chosen treatment.

If there is a full response, continue the treatment. If there is no response or a partial response after adequate dose titration and treatment duration, then go to Stage 2. However, if the patient has a history of comorbid substance abuse, suicidality, or violence, consider moving to Stage 3.

Stage 2

Stage 2 consists of monotherapy with one of the second-generation antipsychotics not tried during Stage 1 or with a first-generation antipsychotic. If the patient achieves full response, continue with the prescribed treatment. If there is no response, or a partial response, after adequate dose titration and treatment duration, go to Stage 3.

Stage 3

Clozapine is the Stage 3 treatment for schizophrenia. Clozapine is the only antipsychotic that has shown superiority over other antipsychotic agents in reducing psychotic symptoms. It is effective in treatment-resistant schizophrenia and in patients having a history of comorbid substance abuse, suicidality, or violence. Persistence of positive symptoms > 2 years warrants and > 5 years requires a clozapine trial, independent of number of preceding antipsychotic trials.

If clozapine does not provide a full response after adequate dose titration and treatment duration, an evaluation should be performed to reexamine the diagnosis and check for substance abuse, medication adherence, and psychosocial stressors. Cognitive behavioral therapy or psychosocial augmentation should also be considered before moving to Stage 4.

Stage 4

Stage 4 treatment includes clozapine plus one of the following:

- A first-generation antipsychotic
- A second-generation antipsychotic
- Electroconvulsive therapy

If there is not a full response after adequate dose titration and treatment duration with any combination pharmacotherapy, move to Stage 5.

Stage 5

Stage 5 treatment consists of monotherapy with a first-generation or second-generation antipsychotic that was not tried at Stages 1 or 2.

The value of Stage 5 treatment in clozapine failures has not been established. If there is partial or no response after adequate dose titration and treatment duration, move to Stage 6.

Stage 6

Stage 6 treatment consists of combination therapy with a second-generation plus a first-generation antipsychotic, a combination of second-generation antipsychotics, a first-generation or second-generation antipsychotic plus electroconvulsive therapy, or a first-generation or second-generation antipsychotic plus another agent such as a mood stabilizer.

The recommendations at Stage 6 are based on case reports, as there have been no controlled studies of combinations in the long-term treatment of schizophrenia.

Tactics and Critical Decision Points

Tactics and Critical Decision Points (CDPs) for the Treatment of Schizophrenia

Stages 1, 2, 4, 5 and 6

Critical Decision Point	Clinical Status	Plan
Week 0 (CDP # 1)	Symptomatic	♦ Assess patient; choose a treatment stage and initiate a medication regimen from that stage; adjust dose to therapeutic dose within one week.
Week 5 (CDP # 2)	Full Response	♦ Continue current dose as maintenance treatment.
	Partial Response	♦ Continue current dose. ♦ Consider increasing dose if medication tolerability is good.
	No Response	♦ Consider the next stage.
Week 8 (CDP # 3)	Full Response	♦ Continue current dose as maintenance treatment.
	Partial Response	♦ Continue current dose. ♦ Consider increasing dose if medication tolerability is good.
	No Response	♦ Consider the next stage.
Week 12 ^{†,§} (CDP # 4)	Full Response	♦ Continue current dose as maintenance treatment.
	Partial Response	♦ Consider the next stage.

[†] Partial responders at week 12 should be titrated to doses at the upper end of the therapeutic range before a change in stages is considered.

[§] ≥ 20% improvement in positive symptoms by week 12 supports continued treatment for another 12 weeks, with dose titration as clinically warranted.

**Tactics and Critical Decision Points (CDPs)
for the Treatment of Schizophrenia**

**Stage 3
Clozapine Monotherapy**

<i>Critical Decision Point</i>	<i>Clinical Status</i>	<i>Plan</i>
Week 0 (CDP # 1)	Symptomatic	<ul style="list-style-type: none"> ◆ Initiate clozapine. Titrate to therapeutic dose within one month. ◆ Adjust dose as needed during interim visits.
Week 16 (CDP # 2)	Full Response	◆ Continue current dose as maintenance treatment.
	Partial Response	◆ Assess serum concentration and adjust dose accordingly.
	No Response	◆ Assess serum concentration and adjust dose accordingly.
Week 28 (CDP # 3)	Full Response	◆ Continue current dose as maintenance treatment.
	Partial Response	<ul style="list-style-type: none"> ◆ Assess serum concentration and adjust dose accordingly. ◆ Consider the next stage.
	No Response	◆ Consider the next stage.

Process Measures: Evaluation of Patient Response

Response

Response criteria are shown below.

RESPONSE CRITERIA

STAGE 1	Positive symptom score ≤ 6
STAGE 2	Positive symptom score ≤ 6
STAGE 3	> 20% decrease in positive symptoms
STAGE 4	> 20% decrease in positive symptoms
STAGE 5	> 20% decrease in positive symptoms
STAGE 6	> 20% decrease in positive symptoms

Generally speaking, symptoms respond to antipsychotics in somewhat different time frames. Agitation, sleep, and appetite often respond during the first 1-2 weeks, whereas personal hygiene and basic interpersonal socialization may be slower to respond (2-3 weeks), and psychotic symptoms can gradually decrease over 2-6 weeks or longer. Residual symptoms may continue to improve at 6-12 weeks. Chronic patients may show slower responses of all symptoms.

Negative symptoms are no longer included in the response criteria, as little evidence exists on which to base realistic goals for negative symptom improvement. Compared to the older agents, the newer medications are thought to be “better” for negative symptoms but this superiority may be explained by the newer agents’ reduced propensity to cause EPS (which can lead to secondary negative symptoms). Several factors (depression, environmental deprivation, positive symptoms) can contribute to negative symptoms, and medications may have little effect on *core* negative symptoms.

This in no way implies that negative symptoms are not important and do not need to be measured. On the contrary, recent findings indicate that negative and cognitive symptoms have more of an impact on patients’ functional status than the positive symptoms of schizophrenia. At each medication visit, clinicians should perform the PSRS, BNSA, and assessments of “other symptoms,” such as mood lability, anxiety, agitation, etc. and incorporate all findings into the clinical decision-making process.

Partial Response

A partial response at any stage of the algorithm is a basis for continuing the patient in that stage, up to the maximum recommended amount of time for that stage. At Critical Decision Points (CDPs), there is the option of changing the antipsychotic dose for partial responders. This is not a requirement, however. For many patients, further duration of treatment may be all that is needed. There are, unfortunately, no empirical guidelines for deciding when this is the case. Generally, prior time to achieve a response in a particular patient is helpful in judging when that patient is likely to

respond to the current treatment. In summary, a partial responder in Stages 1-2 has > 20% improvement in positive symptoms, but his/her absolute positive symptom score exceeds 6. In Stages 3 - 6, partial response is a clinical judgment that the patient whose symptoms have improved by less than 20% is "better." It is not clinically meaningful to try to use scale score changes of less than 20% to distinguish between partial responders and non-responders.

Non-Response

At any stage, before concluding that a patient is a non-responder to an antipsychotic, the clinician should consider causes of non-response that would indicate a course of action other than changing to a new antipsychotic. Included in this list are:

1. Medication non-adherence
2. Incorrect diagnosis
3. Substance abuse
4. Side effects
5. Psychosocial stressors
6. Undiagnosed or uncorrected general medical problem such as diabetes, hypertension, hypothyroidism, and anemia

Process Measures: Evaluation of Patient Response

Positive Symptom Rating Scale (PSRS) and Brief Negative Symptom Assessment (GNSA)

Patients with a diagnosis of schizophrenia or schizoaffective disorder should be evaluated at every visit using the Positive Symptom Rating Scale, or PSRS, and the Brief Negative Symptom Assessment, or BNSA.

PSRS

The 4-item PSRS assesses positive symptoms of schizophrenia (suspiciousness, unusual thought content, hallucinations, and conceptual disorganization). These items are from the Brief Psychiatric Rating Scale (BPRS) (Overall & Gorham, 1962) and the expanded version of the BPRS (Lukoff, et al. 1993), both of which have been shown to be valid and reliable. For the 4-item PSRS the items are ranked on a scale of: N/A = Not Assessed, 1 = Not Present, 2 = Very Mild, 3 = Mild, 4 = Moderate, 5 = Moderately Severe, 6 = Severe, 7 = Extremely Severe. The interview takes 5 minutes or less to complete.

BNSA

The BNSA is a 4-item instrument utilized to assess a subset of DSM-IV negative symptoms (Alogia, Amotivation, Flat Affect, and Asociality). The items are based on items from the Negative Symptom Assessment developed by Alphas et al. (1989) and the Scale for the Assessment of Negative Symptoms (SANS) developed by Andreason (1981). The BNSA provides quick assessment of distinct negative symptoms, and scoring is based largely on observation. The items are scored on a scale of 1 through 6, and the instrument takes less than five minutes to administer.

Clinicians can use the same scoring sheet to record scores from both the PSRS and the BNSA.

A copy of these scales and the scoring sheet can be found in Appendix A.

Clinician Ratings

Each of the symptom clusters is rated on a 10-point scale (from “no symptoms” to “extremely severe”). The rating is based on impression of the patient at this visit, as well as information about the patient’s clinical status during the week prior to the visit.

- **Core Symptoms:** Based upon all available information, clinician impression of the presence and severity of each of the symptoms in this patient.
- **Other Symptoms:** Clinician rating of other symptoms associated with the patient’s disorder, but not core symptoms of the patient’s illness. Rate impressions for each of the specific “other symptoms” listed (irritability, mood lability, insomnia, agitation, anxiety, level of interest, appetite, energy level). Under “other,” specify and rate any other symptoms that are significant.
- **Overall Side Effect Severity:** Overall rating of side effects from all medications being taken by the patient.
- **Overall Functioning:** Overall impression of this patient’s ability to function on a daily basis. “10” is the highest possible functioning, and “1” is the lowest possible functioning.

Patient Self-Ratings

Patient global ratings are recorded on the Clinical Record Form at the beginning of each visit. These ratings should apply to the symptoms and side effects the patient has experienced during the past week, and are rated on a 10-point scale (from “no symptoms” to “extremely severe”).

- **Core Symptoms:** The provider should ask the patient to make a global rating of symptoms he/she has experienced in the past week.
- **Overall Side Effect Severity:** The provider should ask the patient to make a global rating of side effects he/she has experienced in the past week.

Medications and Dosing

Please refer to *Appendix C* for summary of recommended doses, titration schedules, maximum recommended doses, side effects, monitoring parameters, and drug interactions for medication unused in the Algorithm for Treatment of Schizophrenia

Appendix D contains recommendations for dealing with treatment-emergent side effects as well as co-existing symptoms.

Medication Discontinuation

A trial period off antipsychotics may be reasonable for some patients early in the course of illness. This, an individualized decision, depends on a number of factors that do not lend themselves to an algorithmic approach. Although research shows increased relapse rates among patients in discontinuation studies, only minimal guidance is provided regarding this treatment decision in patients who responded well to antipsychotics early in the course of their illness and have maintained a complete remission for a prolonged time period (e.g., > 2 yrs) . Thus, the schizophrenia algorithm contains no guidelines for antipsychotic medication discontinuation, which is anticipated to be a rare event in the typical mental health clinic patient population.

Medication Maintenance

The evidence overwhelmingly favors the conclusion that, for most patients, maintenance antipsychotic medication is a key aspect of successful treatment, in preference to discontinuation or intermittent treatment. Less clear is what the maintenance dose of antipsychotic medication should be for any individual patient. A common clinical aphorism is that the maintenance dose should be the lowest that will keep the patient relatively symptom free. However, very low doses of maintenance medication are clearly less effective for a proportion of patients than doses in the usual range. Moreover, schizophrenia is an illness of natural exacerbations and remissions. Doses that are just sufficient during periods when the illness is quiescent are likely to be inadequate during periods when an exacerbation threatens. That is to say, the optimal maintenance dose is likely to be somewhat higher than the dose which prevents symptoms under the best of circumstances. On the other hand, too high a maintenance dose elevates side effect risks without therapeutic gain.

Documentation

Treatment with the Schizophrenia algorithm utilizes uniform documentation developed by TDSHS and the TMAP team and modified for use by various centers. The critical information from patient history needed for implementation of the SCZ algorithm is:

1. Past and current psychoactive medications and response
2. Primary current diagnosis. (Please note that these algorithms were developed for patients diagnosed with schizophrenia)
3. Core symptoms
4. Other symptoms
5. Side effects (to evaluate tolerability)
6. Response to treatment: overall functioning, PSRS & BNSA scores, patient self-report of symptom severity and side effects

Outpatient Documentation

Forms:

1. Outpatient Clinic Visit Clinical Record Form (CRF): The CRF should be completed at each visit in which a clinician or other clinician is evaluating response to treatment. Please note that all patients will have a stage entered for the principal treatment algorithm.

CRFs may vary in format, but all should contain the minimum data specified *Appendix F*.

2. Process Measures Graphs: The Process Measures Graphs are optional instruments for noting the assessment scores that are recorded at each visit. A new graph is begun if the patient changes algorithm stages. The graphs will give the physician an “at a glance” reference for noting changes or trends in process measures scores as well as for anticipating Critical Decision Points. In order to be useful for decision making, the physician must see the graph at each visit. It can also be useful to show the graph to the patient or family member, thus visually demonstrating the improvement that has occurred since the patient started treatment. Copies of the Process Measures Graphs are included in *Appendix A*.

Optional Forms: If these forms are not used, then an alternative uniform documentation process should be used to record this important information.

1. Outpatient Intake Form
2. Outpatient Interim Contact Form (ICF): In the event that the patient does not come into the clinic or there is not time for a complete visit, the ICF is documented by the physician or other clinical personnel.

3. Patient Algorithm: An individual patient's medication history obtained from patient interview and chart review can be recorded on a "Patient Algorithm", and, when kept up to date, will provide a quick reference for determining a patient's placement in the treatment algorithm.

Inpatient Documentation

Required Forms:

1. Inpatient Clinic Visit Clinical Record Form: Complete as usual. See instructions above for "Outpatient Clinic Visit Clinical Record Form" for detailed example.

Optional Forms:

2. Inpatient Intake Form
3. Inpatient Contact Form

Modifications for Inpatient Use

Patients who have been hospitalized for symptoms of schizophrenia require prompt interventions to achieve stabilization and discharge. It is likely that a clinician may make the following modifications to the TMAP algorithms to achieve these goals.

Adjustment to Critical Decision Points – The algorithm recommends that clinicians see patients every two weeks when a new medication is started, approximately every four weeks while the patient is adjusting to the medication, and no less often than every 3 months once the patient is stable. Of course, opportunities to evaluate the patient and make clinical decisions and medication adjustments happen on an expedited schedule when the patient is an inpatient. Although psychotropic medications do not work faster when a patient is hospitalized, the clinician does have an ongoing opportunity to evaluate the patient's response to treatment. Therefore, critical decision points to evaluate the need for mood stabilizer dosage adjustment or medication change can be made at shorter intervals. In general, if a patient is tolerating usual effective doses of an antipsychotic, dose titration should occur on a weekly basis if needed. Inpatient physicians should fill out a clinical record form for each patient on a weekly basis as well.

Accelerated movement to advanced treatment stage – Admission to a psychiatric unit is almost always due to acute circumstances, such as imminent danger to self or others, grave disablement, and/or a marked exacerbation of symptoms. The necessity of an inpatient admission signals that a change in treatment should be considered and each admission should trigger a thorough evaluation of algorithm staging. Rarely, a patient is admitted for his or her “first break,” and these patients will be started in Stage 1. Far more often the patient has an extensive medication history, and the admitting clinician assumes that the current medication is not working and advances the patient to the next stage of the algorithm. However, before changing a stage, the clinician should evaluate the following four factors:

1. Has the patient been taking the medication? Non-adherence is a major issue in most chronic diseases. Medication does nothing if not taken and, in order to produce maximum benefits, must be taken as directed. Explore this with the patient. Re-starting the current medication may be the best treatment.
2. Is substance abuse a problem? Drug abuse can cause acute and chronic psychiatric symptoms which often remit (albeit slowly) when the abuse stops. Always evaluate for symptoms of withdrawal and, if present, help the patient through the withdrawal period before staging the patient in the algorithm. Keep in mind that patients may resort to drugs of abuse to alleviate medication side effects, especially neurological ones.
3. Is the patient dealing with psychosocial stresses, such as housing problems, family difficulties, and/or employment uncertainties? If so, the treatment team needs to do what it can to help the patient resolve the problem(s), and a change in medication may not be beneficial. However, a medication change is probably warranted if the clinician determines that increased symptomatology was one of the major causes of the patient's psychosocial problem(s).
4. If a patient has not demonstrated at least partial response in positive symptoms after 2-3 weeks treatment, with appropriate dose titration, then the clinician should consider a change in algorithm stage.

Use of adjunctive medications – Patients with schizophrenia frequently have co-existing symptoms of agitation, aggression, excessive anxiety, or insomnia. Inpatient treatment may necessitate the use of adjunctive medications for these symptoms. Refer to Appendix D.

Although it is anticipated that adjunctive medications may be used more commonly in the hospital, their use is still typically time limited, and this intent needs to be communicated to the outpatient treating clinician. For example, at the time of discharge, include instructions for follow-up procedures, including intended taper of short-term medications. Providing the outpatient clinician with the last 1 or 2 inpatient CRFs can be extremely helpful in communicating clinical information.

Inpatient to Outpatient Transition

The transition between inpatient and outpatient care is often unsuccessful. Most inpatient clinicians have dealt with the frustration of discharging a patient only to see him or her return to the hospital within a few weeks as a result of not receiving outpatient follow-up and/or not filling prescriptions. The trend toward brief stays further aggravates the problem by forcing clinicians to discharge patients before they are completely stabilized. By the same token, outpatient clinicians must constantly revise their treatment plans when their long-term treatment intentions are not followed by the inpatient physician. The following three strategies may improve transitions between the two treatment settings:

1. **Document the treatment plan.** It is imperative that all clinicians document the rationale behind treatment decisions and outline the expected treatment plan. This would include detailing expected changes in medications, such as, "I expect Mr. Doe's lorazepam (Ativan®) for agitation to be discontinued once agitation has either subsided or is adequately controlled with continued treatment of his antipsychotic medication (olanzapine)." This intent needs to be communicated to outpatient treating clinicians.

Use of uniform documentation among all clinicians involved in a patient's care assists in addressing this, and the TMAP documentation system is an example of such an effort. Additionally, inpatient clinicians may want to start notes to their outpatient colleagues with "transfer" rather than "discharge" (I am 'transferring' the acute care of this patient...) because the former term implies a continuation of care while the latter suggests a disruption.

Similarly, it is important that clinicians treating patients in the hospital receive documentation of the care provided in the clinic in a timely fashion. Delays in the communication of outpatient treatment information to the inpatient physician may result in inappropriate treatment regimens and prolonged times to symptom stabilization and discharge. Once again, the use of uniform documentation systems can facilitate this process.

2. **Ensure that patients leave the hospital with enough medication to see them through to the first follow-up appointment.** If administrative policies prevent adequate supplies of medication from being dispensed, these policies need to be challenged.
3. **Establish communication between the inpatient and outpatient treatment teams.** Physicians working in both the inpatient and outpatient arenas need to get to know one another and brainstorm methods to improve coordination of care between the two settings. Three possible strategies for improving communication are:
 - i. Have a team member (on each side) coordinate and follow-up on transfers.
 - ii. Organize quarterly meetings with key inpatient and outpatient staff members.
 - iii. Designate individuals in the respective organizations responsible for communicating patient information to clinicians in the other treatment setting. In the clinic, this may include methods to communicate information to hospital clinicians during the night and on weekends.
4. **Use of clinical report form (CRF):** If the clinician documents pharmacotherapy care on the CRF, then a transfer of copies of the last 1 or 2 completed CRF's to the clinician assuming care of the patient can be helpful in communicating the treatment the patient has received as well as the clinical status the last time the patient was seen.

Outpatient to Inpatient Transition

Communication and transition in care is equally important when a patient is admitted to the hospital. The outpatient treating clinician should be contacted when patients are hospitalized, and copies of the last two CRFs should be faxed to the hospital. The outpatient clinician should be asked about the patient's response to medication and potential reasons for illness exacerbation. It should not necessarily be assumed that a patient relapsed because of medication treatment failure. Not taking medications appropriately and alcohol or other substance use are common factors leading to hospitalization. These as well as other factors (e.g., family or other environmental stress) should be considered in deciding whether to continue the patient on the same medication regimen being used in the outpatient setting or to move to a new treatment stage.

Appendix A: Process Measures

- Administration Manual, 4-item Positive Symptom Rating Scale (PSRS) and Brief Negative Symptom Assessment (BNSA)
- PSRS and BNSA Score Sheet
- Process Measures Graphs
- Scoring Criteria for Clinician- and Patient-Rated Overall Symptom and Side Effect Ratings

4-ITEM POSITIVE SYMPTOM RATING SCALE (PSRS) (Version 5.0)

SCALE ITEMS AND ANCHOR POINTS

1. **SUSPICIOUSNESS:** Expressed or apparent belief that other persons have acted maliciously or with discriminatory intent. Include persecution by supernatural or other nonhuman agencies (e.g., the devil). Note: Ratings of “3” or above should also be rated under Unusual Thought Content.

*Do you ever feel uncomfortable in public? Does it seem as though others are watching you?
Are you concerned about anyone's intentions toward you?
Is anyone going out of their way to give you a hard time, or trying to hurt you? Do you feel in any danger?*

[If patient reports any persecutory ideas/delusions, ask the following]:

How often have you been concerned that [use patient's description]? Have you told anyone about these experiences?

1 Not Present

2 Very Mild

Seems on guard. Reluctant to respond to some “personal” questions. Reports being overly self-conscious in public.

3 Mild

Describes incidents in which others have harmed or wanted to harm him/her that sound plausible. Patient feels as if others are watching, laughing, or criticizing him/her in public, but this occurs only occasionally or rarely. Little or no preoccupation.

4 Moderate

Says others are talking about him/her maliciously, have negative intentions, or may harm him/her. Beyond the likelihood of plausibility, but not delusional. Incidents of suspected persecution occur occasionally (less than once per week) with some preoccupation.

5 Moderately Severe

Same as 4, but incidents occur frequently, such as more than once per week. Patient is moderately preoccupied with ideas of persecution OR patient reports persecutory delusions expressed with much doubt (e.g., partial delusion).

6 Severe

Delusional -- speaks of Mafia plots, the FBI, or others poisoning his/her food, persecution by supernatural forces.

7 Extremely Severe

Same as 6, but the beliefs are bizarre or more preoccupying. Patient tends to disclose or act on persecutory delusions.

- 2. UNUSUAL THOUGHT CONTENT:** Unusual, odd, strange or bizarre thought content. Rate the degree of unusualness, not the degree of disorganization of speech. Delusions are patently absurd, clearly false or bizarre ideas that are expressed with full conviction. Consider the patient to have full conviction if he/she has acted as though the delusional belief were true. Ideas of reference/persecution can be differentiated from delusions in that ideas are expressed with much doubt and contain more elements of reality. Include thought insertion, withdrawal and broadcast. Include grandiose, somatic and persecutory delusions even if rated elsewhere. Note: If Suspiciousness is rated "6" or "7" due to delusions, then Unusual Thought Content must be rated a "4" or above.

Have you been receiving any special messages from people or from the way things are arranged around you? Have you seen any references to yourself on TV or in the newspapers?

Can anyone read your mind?

Do you have a special relationship with God?

Is anything like electricity, X-rays, or radio waves affecting you?

Are thoughts put into your head that are not your own?

Have you felt that you were under the control of another person or force?

[If patient reports any odd ideas/delusions, ask the following]:

How often do you think about [use patient's description]?

Have you told anyone about these experiences? How do you explain the things that have been happening [specify]?

1 Not Present

2 Very Mild

Ideas of reference (people may stare or may laugh at him), ideas of persecution (people may mistreat him). Unusual beliefs in psychic powers, spirits, UFOs, or unrealistic beliefs in one's own abilities. Not strongly held. Some doubt.

3 Mild

Same as 2, but degree of reality distortion is more severe as indicated by highly unusual ideas or greater conviction. Content may be typical of delusions (even bizarre), but without full conviction. The delusion does not seem to have fully formed, but is considered as one possible explanation for an unusual experience.

4 Moderate

Delusion present but no preoccupation or functional impairment. May be an encapsulated delusion or a firmly endorsed absurd belief about past delusional circumstances.

5 Moderately Severe

Full delusion(s) present with some preoccupation OR some areas of functioning disrupted by delusional thinking.

6 Severe

Full delusion(s) present with much preoccupation OR many areas of functioning are disrupted by delusional thinking.

7 Extremely Severe

Full delusions present with almost total preoccupation OR most areas of functioning are disrupted by delusional thinking.

In the past 7 days...

3. **HALLUCINATIONS:** Reports of perceptual experiences in the absence of relevant external stimuli. When rating degree to which functioning is disrupted by hallucinations, include preoccupation with the content and experience of the hallucinations, as well as functioning disrupted by acting out on the hallucinatory content (e.g., engaging in deviant behavior due to command hallucinations). Include "thoughts aloud" ("gedankenlautwerden") or pseudohallucinations (e.g., hears a voice inside head) if a voice quality is present.

Do you ever seem to hear your name being called?

Have you heard any sounds or people talking to you or about you when there has been nobody around?

[If hears voices]: What does the voice/voices say? Did it have a voice quality?

Do you ever have visions or see things that others do not see? What about smell — odors that others do not smell?

[If the patient reports hallucinations, ask the following]:

Have these experiences interfered with your ability to perform your usual activities/work?

How do you explain them? How often do they occur?

1 Not Present

2 Very Mild

While resting or going to sleep, sees visions, smells odors, or hears voices, sounds or whispers in the absence of external stimulation, but no impairment in functioning.

3 Mild

While in a clear state of consciousness, hears a voice calling the subject's name, experiences non-verbal auditory hallucinations (e.g., sounds or whispers), formless visual hallucinations, or has sensory experiences in the presence of a modality-relevant stimulus (e.g., visual illusions) infrequently (e.g., 1-2 times per week) and with no functional impairment.

4 Moderate

Occasional verbal, visual, gustatory, olfactory, or tactile hallucinations with no functional impairment OR non-verbal auditory hallucinations/visual illusions more than infrequently or with impairment.

5 Moderately Severe

Experiences daily hallucinations OR some areas of functioning are disrupted by hallucinations.

6 Severe

Experiences verbal or visual hallucinations several times a day OR many areas of functioning are disrupted by these hallucinations.

7 Extremely Severe

Persistent verbal or visual hallucinations throughout the day OR most areas of functioning are disrupted by these hallucinations.

4. CONCEPTUAL DISORGANIZATION: Degree to which speech is confused, disconnected, vague or disorganized. Rate tangentiality, circumstantiality, sudden topic shifts, incoherence, derailment, blocking, neologisms, and other speech disorders. Do not rate content of speech.

1 Not Present

2 Very Mild

Peculiar use of words or rambling but speech is comprehensible.

3 Mild

Speech a bit hard to understand or make sense of due to tangentiality, circumstantiality, or sudden topic shifts.

4 Moderate

Speech difficult to understand due to tangentiality, circumstantiality, idiosyncratic speech, or topic shifts on many occasions OR 1-2 instances of incoherent phrases.

5 Moderately Severe

Speech difficult to understand due to circumstantiality, tangentiality, neologisms, blocking, or topic shifts most of the time OR 3-5 instances of incoherent phrases.

6 Severe

Speech is incomprehensible due to severe impairments most of the time. Many PSRS items cannot be rated by self-report alone.

7 Extremely Severe

Speech is incomprehensible throughout interview.

Sources of information (check all applicable):

- Patient
- Parents/Relatives
- Mental Health Professionals
- Chart

Explain here if validity of assessment is questionable:

- Symptoms possibly drug-induced
- Underreported due to lack of rapport
- Underreported due to negative symptoms
- Patient uncooperative
- Difficult to assess due to formal thought disorder
- Other _____

Confidence in assessment:

_____ 1 = Not at all - 5 = Very confident

BRIEF NEGATIVE SYMPTOM ASSESSMENT SCALE (version 5.0)

Items adapted from NSA and SANS

1. **PROLONGED TIME TO RESPOND** (a measure of Alogia): Observed throughout communication with the patient. After asking the patient a question, he or she pauses for inappropriately long periods before initiating a response. Delay is considered a pause if it feels as though you are waiting for a response or if you consider repeating the question because it appears that the patient has not heard you. He or she may seem “distant,” and sometimes the examiner may wonder if he has even heard the question. Prompting usually indicates that the patient is aware of the question, but has been having difficulty in developing his thoughts in order to make an appropriate reply. Rate severity on the frequency of these pauses.
 - 1 **Normal**
No abnormal pauses before speaking.
 - 2 **Minimal**
Minimal evidence of inappropriate pauses (brief but not abnormally lengthy pauses occur) may be extreme of normal
 - 3 **Mild**
Occasional noticeable pauses before answering questions. Due to the length of the pause, you feel the need to repeat yourself once or twice during the interview.
 - 4 **Moderate**
Distinct pauses occur frequently (20-40% of responses).
 - 5 **Marked**
Distinct pauses occur most of the time (40-80% of responses).
 - 6 **Severe**
Distinct pauses occur with almost every response (80-100% of responses).

2. **EMOTION: UNCHANGING FACIAL EXPRESSION; BLANK, EXPRESSIONLESS FACE** (a measure of Flat Affect): The patient’s face appears wooden, mechanical, frozen. Facial musculature is generally expressionless and unchanging. The patient does not change expression, or change is less than normally expected, as the emotional content of discourse changes. Because of this, emotions may be difficult to infer. Disregard changes in facial expression due to abnormal involuntary movements, such as tics and tardive dyskinesia. The two dimensions of importance when making this rating are degree of emotional expression and spontaneity.
 1. **Normal**
Spontaneous displays of emotion occur when expected. Normal degree of expressiveness of emotions is present.
 2. **Minimal**
Spontaneous expressions of emotion occur when expected. However, there is a reduction in degree or intensity of the emotions expressed. May be extreme of normal.
 3. **Mild**
Spontaneous expressions of emotion occur infrequently. When emotions are expressed, there is a reduction in degree or intensity displayed.

4. **Moderate**
Obvious reduction in spontaneous expressions. Spontaneous expressions of emotion may occur very rarely during interaction and only when discussing topics of special interest or humor to the subject.
5. **Marked**
Facial expression is markedly decreased. There are no spontaneous expressions of emotion unless prompted or coaxed by the interviewer.
6. **Severe**
There are no expressions of emotion even when attempts are made to elicit an emotional response. The subject's face remains blank throughout the interview.

3. REDUCED SOCIAL DRIVE. (a measure of Asociality): This item assesses how much the subject desires to initiate social interactions. Desire may be measured in part by the number of actual or attempted social contacts with others. If the patient has frequent contact with someone (e.g., family member) who initiates the contact, does the patient appear to desire the contact (i.e., would he or she initiate contact if necessary)? In making this rating, probe the desire to initiate social interactions, number of social interactions, and the ability to enjoy them.

Assessed by asking the patient questions like:

How have you spent your time in the past week?

Do you live alone or with someone else?

Do you like to be around people?

Do you spend much time with others?

Do you have difficulty feeling close to others?

Who are your friends?

How often do you see them?

Did you see them this past week?

Have you called them on the phone?

When you get together, who decides what to do and where to go?

When you spend time with others, do you ask them to do something with you or do you wait until they ask you to do something?

Is anyone concerned about your happiness or well being?

- 1 **Normal**
Normal desire to initiate and normal number of contacts. Social contacts are enjoyable.
- 2 **Minimal**
Minimal reduction in either the desire to initiate social contacts or the number of social relationships. May initially seem guarded, but has the ability to establish relationships over time. Social relationships are enjoyable.
- 3 **Mild**
Reduction in desire to initiate social contacts. The patient has few social relationships and these social contacts are enjoyable.
- 4 **Moderate**
Obvious reduction in the desire to initiate social contacts. The patient has few relationships toward which he or she feels indifference. However, a number of social contacts are initiated each week.

- 5 Marked**
Marked reduction in desire to initiate social contacts. The patient has very few relationships toward which he or she feels indifference. The patient does not initiate social contacts but may maintain a few contacts (such as with family).
- 6 Severe**
Patient does not desire social contact. Actively avoids social interactions.

- 4. GROOMING AND HYGIENE** (a measure of Amotivation): Observed during interaction with the patient. The patient displays less attention to grooming and hygiene than normal. The patient presents with poorly groomed hair, disheveled clothing, etc. Do not rate grooming as poor if it is simply done in what one might consider poor taste (e.g., wild hairdo or excessive makeup). In addition to observation, one must ask the patient about regularity of bathing, brushing teeth, changing clothes, etc. This is particularly important with outpatients, as the patient may present his or her best grooming and hygiene at their clinic visit. Two dimensions to keep in mind when making this rating are current appearance and regularity of grooming behaviors.

Assess the patient by asking questions like:

*How many times in the past week have you taken a shower or bath?
How often do you change your clothes?
How often do you shower and brush your teeth?*

- 1 Normal**
Patient is clean (e.g., showers every day) and dressed neatly.
- 2 Minimal**
Minimal reduction in grooming and hygiene, may be at the extreme end of the normal range.
- 3 Mild**
Apparently clean but untidy appearance. Clothing may be mismatched. Patient may shower less often than every other day, or may brush teeth less than everyday.
- 4 Moderate**
There is an obvious reduction in grooming and hygiene. Clothes may appear unkempt, rumpled, or the patient may look as if he or she just got out of bed. The patient may go without showering or bathing for two days at a time. The patient may go for two days without brushing his/her teeth.
- 5 Marked**
There is a marked reduction in grooming and hygiene. Clothing may appear dirty, stained, or very unkempt. The subject may have greasy hair or a body odor. The patient may go 3 days at a time without showering or 3 or 4 days without brushing his/her teeth.
- 6 Severe**
Clothing is badly soiled. Patient has a foul odor. Patient may go more than 4 days in a row without showering or more than 4 days in a row without brushing his/her teeth. Poor hygiene may present a health risk.

SCORE SHEET for 4-ITEM POSITIVE SYMPTOM RATING SCALE AND BRIEF NEGATIVE SYMPTOM ASSESSMENT

4-Item Positive Symptom Rating Scale

Use each item's anchor points to rate the patient.

1. Suspiciousness	*NA	1	2	3	4	5	6	7	
2. Unusual Thought Content	NA	1	2	3	4	5	6	7	
3. Hallucinations	NA	1	2	3	4	5	6	7	
4. Conceptual Disorganization	NA	1	2	3	4	5	6	7	SCORE: _____

* NA – not able to be assessed

4-Item Negative Symptom Rating Scale

Use each item's anchor points to rate the patient.

1. Prolonged Time to Respond	1	2	3	4	5	6	
2. Emotion Unchanging facial expression, blank, expressionless face.	1	2	3	4	5	6	
3. Reduced Social Drive	1	2	3	4	5	6	
4. Poor Grooming and Hygiene	1	2	3	4	5	6	SCORE: _____

Source of Information (check all applicable)

Explain here if validity of assessment is questionable:

- Patient
- Parents/Relatives
- Mental Health Professionals
- Chart

- Symptoms possibly drug-induced
- Underreported due to lack of rapport
- Underreported due to negative symptoms
- Patient uncooperative
- Difficult to assess due to formal thought disorder
- Other _____

Confidence in assessment
___ 1=Not at all – 5=Very confident

The 4-item PSRS was adapted from the Expanded Version of the BPRS developed by:
Ventura, J.; Lukoff, D.; Nuechterlein, K.H.; Liberman, R.P.; Green, M.F.; and Shaner, A. Manual for the expanded Brief Psychiatric Rating Scale. *International Journal of Methods Psychiatry Research*, 3:227-244, 1993

The Brief Negative Symptom Assessment was adapted from the Negative Symptom Assessment and the Scale for the Assessment of Negative Symptoms developed respectively by:

Alphs and Summerfelt. *The Negative Symptom Assessment: A new instrument to assess negative symptoms of schizophrenia*. Psychopharmacology Bulletin, 1989. 25(2): p. 159-163.
Andreason, N. *Modified scale for the assessment of negative symptoms. NIMH treatment strategies in schizophrenia study*. Public Health Administration U.S. Department of Health and Human Services, 1984. ADM (9/85): p. 9-102.

Appendix B: Communications

TMAP Information

The University of Texas at Austin
College of Pharmacy PHR 5.110
1 University Station A1910
Austin, TX 78712

TMAP Phone: 512-232-5986
TMAP Fax: 512-735-TMAP (8627)
TMAP Email: info@WebTMAP.org

Appendix C: Medication Charts

Medications Included in Algorithms for Most Recent Episode of Schizophrenia

(Please refer to the Physicians' Desk Reference, FDA approved product labeling, or other sources for more complete information.)

<i>Antipsychotics, Second Generation</i>	<i>40</i>
<i>Antipsychotics, First Generation.....</i>	<i>42</i>
<i>Adjunctive Agents, Agitation/Insomnia</i>	<i>44</i>
<i>Adjunctive Agents, Insomnia.....</i>	<i>44</i>
<i>Adjunctive Agents, Aggression/Hostility.....</i>	<i>45</i>
<i>Adjunctive Agents, Depression</i>	<i>47</i>
<i>Additional References for Drug Information</i>	<i>51</i>

Appendix C: Medication Charts

Antipsychotics, Second Generation								
Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Aripiprazole Abilify®	10-15 mg/day	None	10-30 mg/day	30 mg/day	Once daily	1) Pregnancy test – as clinically indicated 2) BMI measurement – when a new antipsychotic is initiated, at every visit (monthly for inpatients) for 6 months after the new antipsychotic is initiated and quarterly when the antipsychotic dose is stable. 3) Fasting plasma glucose level or hemoglobin A1c – before initiating a new antipsychotic, then yearly. If a patient has significant risk factors for diabetes and for those that are gaining weight – before initiating a new antipsychotic, 4 months after starting an antipsychotic, and then yearly.	<ul style="list-style-type: none"> • Agitation • Constipation • EPS • Insomnia • Nausea • Somnolence 	<ul style="list-style-type: none"> • Carbamazepine • Fluoxetine • Ketoconazole • Paroxetine • Quinidine • St John's Wort
Olanzapine Zyprexa®	5-10 mg/day	5 mg/wk	10-20 mg/day	30 mg/day ²	Once daily	4) Lipid screening [total cholesterol, low-and high-density lipoprotein (LDL and HDL) cholesterol, and triglycerides] – Every 2 years or more often if lipid levels are in the normal range, every 6 months if the LDL level is > 130 mg/dl 5) EKG – before initiating treatment with ziprasidone (Geodon ®) and subsequently if the patient demonstrates symptoms (e.g., syncope) associated with QT interval prolongation.	<ul style="list-style-type: none"> • Constipation • Dizziness • Dry mouth • Glucose dysregulation • Hyperlipidemia • Increased appetite • Sedation • Weight gain 	<ul style="list-style-type: none"> • Carbamazepine • Fluvoxamine • Rifampin • Smoking • St. John's Wort
Quetiapine Seroquel®	50 mg/day	300 mg every 3-7 days	300-800 mg/day	800 mg/day	Twice daily	6) Sexual function inquiry – inquire yearly for evidence of galactorrhea/gynecomastia, menstrual disturbance, libido disturbance or erectile/ejaculatory disturbances in males. If a patient is receiving an antipsychotic known to be associated with Prolactin elevation, then at each visit (quarterly for inpatients) for the first 12 months after starting an antipsychotic or until the medication dose is stable and then yearly. 7) Prolactin level – if there is evidence of galactorrhea/gynecomastia, menstrual disturbance, libido disturbance or erectile/ejaculatory disturbances in males.	<ul style="list-style-type: none"> • Cataract formation • Dry mouth • Glucose dysregulation • Headache • Hyperlipidemia • Increased appetite • Orthostatic hypotension • Sedation • Weight gain 	<ul style="list-style-type: none"> • Erythromycin • Fluconazole • Ketoconazole • Phenytoin • St. John's Wort • Thioridazine • Valproate
Risperidone Risperdal®	1-2 mg/day	1 mg every 2-3 days	2-6 mg/day	16 mg/day ³	Once daily	8) EPS Evaluation (examination for rigidity, tremor, akathisia) – before initiation of any antipsychotic medication, then weekly for the first 2 weeks after initiating treatment with a new antipsychotic or until the dose has been stabilized and weekly for 2 weeks after a dose increase 9) Tardive dyskinesia evaluation – every 6 months. For high-risk patients (including the elderly) every 3 months. 10) Vision questionnaire – ask whether the patient has experienced a change in vision and should specifically ask about distance vision and blurry vision – yearly. 11) Ocular evaluations – yearly for patients older than age 40 years; every 2 years for younger patients	<ul style="list-style-type: none"> • EPS • Glucose dysregulation • Galactorrhea • Hyperlipidemia • Menstrual irregularity • Orthostatic hypotension • Prolactin elevation • Sedation • Sexual dysfunction • Tardive dyskinesia • Weight gain 	<ul style="list-style-type: none"> • Carbamazepine • Cimetidine • Fluoxetine • Paroxetine • Phenytoin • Rifampin • Tricyclic antidepressants

² The CATIE trial utilized olanzapine doses up to 30 mg/day.

³ Use of risperidone > 6mg/day is associated with an increased risk of EPS.

Appendix C: Medication Charts

Antipsychotics, Second Generation								
Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Ziprasidone Geodon®	40- 80 mg/day	20-40 mg every 2-3 days	80 - 160 mg/day	160 mg/day	Once or twice daily ⁴	See Previous Page	<ul style="list-style-type: none"> • Dizziness • ECG changes • EPS • Rash • Sedation • Vomiting 	<ul style="list-style-type: none"> • Carbamazepine • Diuretics • Moxifloxacin • Quinidine • Sotalol • Thioridazine • Tricyclic antidepressants
Clozapine <i>Generic available</i> Clozaril® Fazaclo®	12.5 mg/day (half of a 25 mg tab)	<p>Day 2: 25 mg daily Day 3: 25 mg twice daily Day 6: 25 mg at morning, 50 mg at evening Day 9: 50 mg twice daily Day 12: 75 mg twice daily Day 15: 100 mg twice daily Day 18: 125 mg twice daily Day 21: 150 mg twice daily Day 24: 100 mg at morning, 200 mg at evening</p>	300-900 mg/day (serum level for doses>600 mg/day)	900 mg/day	Twice daily ⁵	<p>1) CBC as indicated by guidelines approved by the FDA in the product labeling.</p> <p>2) Pregnancy test – as clinically indicated</p> <p>3) BMI measurement – when a new antipsychotic is initiated, at every visit (monthly for inpatients) for 6 months after the new antipsychotic is initiated, and quarterly when the antipsychotic does is stable.</p> <p>4) Fasting plasma glucose level or hemoglobin A1c - before initiating a new antipsychotic, then yearly. If a patient has significant risk factors for diabetes and for those that are gaining weight – before initiating a new antipsychotic, 4 months after starting an antipsychotic, and then yearly.</p> <p>5) Lipid screening [total cholesterol, low-and high-density lipoprotein (LDL and HDL) cholesterol, and triglycerides] – Every 2 years or more often if lipid levels are in the normal range, every 6 months if the LDL level is > 130 mg/dl</p> <p>6) Sexual function inquiry – inquire yearly for evidence of galactorrhea/gynecomastia, menstrual disturbance, libido disturbance or erectile/ejaculatory disturbances in males. If a patient is receiving an antipsychotic known to be associated with Prolactin elevation, then at each visit (quarterly for inpatients) for the first 12 months after starting an antipsychotic or until the medication dose is stable and then yearly.</p> <p>7) EPS Evaluation (examination for rigidity, tremor, akathisia) – before initiation of any antipsychotic medication, then weekly for the first 2 weeks after initiating treatment with a new antipsychotic or until the dose has been stabilized and weekly for 2 weeks after a dose increase.</p> <p>8) Tardive Dyskinesia evaluation – every 12 months. For high risk patients (including the elderly), every 6 months.</p> <p>9) Vision questionnaire – ask whether the patient has experienced a change in vision and should specifically ask about distance vision and blurry vision – yearly</p> <p>10) Ocular evaluations – yearly for patients older than age 40 years; every 2 years for younger patients</p>	<ul style="list-style-type: none"> • Agranulocytosis • Excess salivation • Fever • Glucose dysregulation • Hyperlipidemia • Increased appetite • Myocarditis • Orthostatic hypotension • Sedation • Seizures • Tachycardia • Weight gain 	<ul style="list-style-type: none"> • Barbiturates • Caffeine • Carbamazepine • Cimetidine • Erythromycin • Phenytoin • Rifampin • Ritonavir • Smoking • SSRIs • St John's Wort

⁴ The presence of food can increase ziprasidone's absorption up to two-fold.

⁵ Eventual maintenance dose schedule is twice daily dosing. 1/3 dose in morning, 2/3 dose in the evening.

Appendix C: Medication Charts

Antipsychotics, First Generation

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Low Potency						<p>1) Pregnancy test – as clinically indicated</p> <p>2) BMI measurement – when a new antipsychotic is initiated, at every visit (monthly for inpatients) for 6 months after the new antipsychotic is initiated, and quarterly when the antipsychotic dose is stable</p> <p>3) Fasting plasma glucose level or hemoglobin A1c – before initiating a new antipsychotic, then yearly</p> <p>If a patient has significant risk factors for diabetes and for those that are gaining weight – before initiating a new antipsychotic, 4 months after starting an antipsychotic, and then yearly</p> <p>4) Lipid screening [total cholesterol, low-and high-density lipoprotein (LDL and HDL) cholesterol, and triglycerides] – Every 2 years or more often if lipid levels are in the normal range, every 6 months if the LDL level is > 130 mg/dl</p> <p>5) Sexual function inquiry – inquire yearly for evidence of galactorrhea/gynecomastia, menstrual disturbance, libido disturbance or erectile/ejaculatory disturbances in males.</p> <p>If a patient is receiving an antipsychotic known to be associated with prolactin elevation, then at each visit (quarterly for inpatients) for the first 12 months after starting an antipsychotic or until the medication dose is stable and then yearly</p> <p>6) Prolactin level – if there is evidence of galactorrhea/gynecomastia, menstrual disturbance, libido disturbance or erectile/ejaculatory disturbances in males.</p> <p>7) EPS Evaluation (examination for rigidity, tremor, akathisia) – before initiation of any antipsychotic medication, then weekly for the first 2 weeks after initiating treatment with a new antipsychotic or until the dose has been stabilized and weekly for 2 weeks after a dose increase</p> <p>8) Tardive dyskinesia evaluation – every 6 months</p> <p>For high risk patients (including the elderly), every 3 months.</p> <p style="text-align: center;">Continued on next page</p>	<ul style="list-style-type: none"> • Constipation • Dry mouth • EPS • Orthostatic hypotension • Photosensitivity • Sedation • Tachycardia • Tardive dyskinesia <p style="text-align: center;">See Previous Page</p>	<ul style="list-style-type: none"> • Guanethidine • Meperidine • Paroxetine • Pindolol • Quinolones • β-Blockers • Ziprasidone
Mid Potency								<ul style="list-style-type: none"> • Barbiturates • General anesthetics • Opiate agonists • Radiopaque contrast agents • Tricyclic antidepressants
Chlorpromazine <i>Generic available</i> Thorazine®	50-100 mg/day	50-200 mg/day	300-1000 mg/day	1000 mg/day	Three times daily			
Loxapine <i>Generic available</i> Loxitane®	20 mg/day	10-50 mg every 3-7 days	50-150 mg/day	150 mg/day	Once or twice daily			
Molindone Moban®	50-75 mg/day	25 mg every 3 days	50-150 mg/day	225 mg/day	3-4 times daily			<ul style="list-style-type: none"> • CNS depressants • Anticonvulsants • Tetracycline
Perphenazine <i>Generic available</i> Trilafon®	4-8 mg/day	4-8 mg/day	16-64 mg/day	64 mg/day ⁶	Three times daily	<ul style="list-style-type: none"> • Paroxetine • Quinolones 		
High Potency						<ul style="list-style-type: none"> • Guanethidine • Paroxetine • Quinolones 		
Fluphenazine <i>Generic available</i> Prolixin®	5 mg	2.5-5 mg/day	5-20 mg/day	20 mg/day	Three times daily			
Fluphenazine D <i>Generic available</i> Prolixin Decanoate®	12.5-25 mg IM every 1-3 weeks ⁷	12.5 mg per injection	6.25-50 mg IM every 2-4 weeks	100mg IM (per 4 weeks)	Every 1-3 weeks	<ul style="list-style-type: none"> • Guanethidine • Paroxetine • Quinolones 		

⁶ The CATIE trial used doses up to 32mg/day.

⁷ Starting dose generally 1.2 times the patient's oral dose.

Appendix C: Medication Charts

Antipsychotics, First Generation

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Haloperidol Generic available Haldol®	2-5 mg/day	2-5 mg/day	2-20 mg/day	20 mg/day	1 - 3 times daily	9) Vision questionnaire –ask whether the patient has experienced a change in vision and should specifically ask about distance vision and blurry vision – yearly 10) Ocular evaluations – yearly for patients older than age 40 years; every 2 years for younger patients	See previous page	<ul style="list-style-type: none"> • Azole antifungals • Carbamazepine • Rifabutin • Rifampin
Haloperidol D Generic available Haldol Decanoate®	25-50 mg IM every 2-4 weeks ⁸	N/A	50-200 mg IM every 2-4 weeks	450 mg (per 4 weeks) ^{9,10}	Every 3-4 weeks			
Thiothixene Navane®	5-10 mg/day	5-10 mg every 3-7 days	15-50 mg/day	50 mg/day	2-3 times daily			<ul style="list-style-type: none"> • Guanethidine
Trifluoperazine Stelazine®	2-5 mg twice daily	2-5 mg every 3-7 days	5-20 mg/day	40 mg/day	Twice daily			<ul style="list-style-type: none"> • Alcohol • Cisapride • Guanethidine • Metrizamide • Paroxetine

⁸ Starting dose generally 10-20 times the patient's oral dose. Dose of first injection should not exceed 100 mg.

⁹ The maximum volume per injection site should not exceed 3 mL.

¹⁰ Multiple injections can be given at 1-7 day intervals to provide total loading dose.

Appendix C: Medication Charts

Adjunctive Agents, Agitation/Insomnia

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Clonazepam <i>Generic Available</i> Klonopin®	0.25-2 mg twice daily	N/A	0.5-4 mg/day	20 mg/day	As needed	1) Pregnancy Test – as clinically indicated	<ul style="list-style-type: none"> • Dizziness • Sedation • Dyskinesia • Slurred speech 	<ul style="list-style-type: none"> • Sodium oxybate • Phenytoin • Carbamazepine • Azole antifungals
Lorazepam <i>Generic Available</i> Ativan®	0.5-2 mg Three times daily	N/A	1-8 mg/day	10 mg/day	As needed			<ul style="list-style-type: none"> • Aminophylline • Clozapine • Kava • Probenecid • Sodium oxybate • Theophylline • Valproate
Temazepam <i>Generic Available</i> Restoril®	7.5-30 mg at bedtime	N/A	15-30 mg/day	30 mg/day	Once daily at bedtime			<ul style="list-style-type: none"> • CNS depressants • Kava • Valerian

Adjunctive Agents, Insomnia

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Eszopiclone Lunesta®	2-3 mg	N/A	2-3 mg/day	3 mg/day	Once daily at bedtime	1) Pregnancy Test – as clinically indicated	<ul style="list-style-type: none"> • Altered taste • Headache • Dry mouth • Dizziness 	<ul style="list-style-type: none"> • Pramipexole • St. John's Wort • CNS depressants • Kava • Valerian
Trazodone <i>Generic available</i> Desyrel®	25 mg	N/A	12.5 – 100 mg/day	400 mg/day	Once daily at bedtime	1) Pregnancy Test – as clinically indicated 2) EKG – as clinically indicated	<ul style="list-style-type: none"> • Blurred vision • Dizziness • Orthostatic hypotension • Dry mouth • Priapism 	<ul style="list-style-type: none"> • Digoxin • Anticonvulsants • Antihistamines • Sedatives • Tranquilizers • Barbiturates • Protease inhibitors • Anticoagulants • SSRIs
Zolpidem <i>Generic available</i> Ambien®	5 mg	N/A	5-10 mg/day	10 mg/day	Once daily at bedtime	1) Pregnancy Test – as clinically indicated	<ul style="list-style-type: none"> • Dizziness • Somnambulism • Confusion • Headache 	<ul style="list-style-type: none"> • Protease inhibitors • Pramipexole • CNS depressants • Kava • Valerian
Zaleplon Sonata®	5-10 mg	N/A	5-10 mg/day	20 mg/day	Once daily at bedtime		<ul style="list-style-type: none"> • Dizziness • Abdominal pain • Headache • Paresthesia 	<ul style="list-style-type: none"> • Cimetidine • Phenobarbital • Carbamazepine • Protease inhibitors • CNS depressants • Kava • Valerian

N/A = Not Applicable

Appendix C: Medication Charts

Adjunctive Agents, Aggression/Hostility

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Carbamazepine <i>Generic available</i> Tegretol® Tegretol® XR Carbatrol®	200-600 mg/day	200 mg/day every 2-4 days	400-1600 mg/day	1600 mg/day ^{11,12}	2 - 3 Times Daily	1) CBC with platelets – baseline and 1 to 2 weeks after each dose increase and as clinically indicated 2) Hepatic function panel and electrolytes; baseline and as clinically indicated 3) Pregnancy Test – as clinically indicated 4) Carbamazepine Levels – 1-2 weeks after dose adjustment, then as clinically indicated Therapeutic Serum Concentration ¹³ : 4-12 mcg/ml Frequency of Serum Level Monitoring: <ul style="list-style-type: none"> • Day 5-7 • Weekly until stable • Every 3-6 months 	<ul style="list-style-type: none"> • Ataxia • Diplopia • Dizziness • Dysarthria • GI upset • Hyponatremia • Leukopenia • Nystagmus • Rash • Sedation 	<ul style="list-style-type: none"> • Antipsychotics • Benzodiazepines • Cimetidine • Corticosteroids • Divalproex • Erythromycin • Lamotrigine¹² • Oral contraceptive pill • SSRIs • Tricyclic antidepressants • Warfarin • Induce its own metabolism. May require close dose titration.
Valproic Acid, Divalproex Sodium <i>Generic available</i> Depakene® Depakote®	500 - 1000 mg/day	500 mg/day every 1-2 weeks	750-2000 mg/day	60 mg/kg/day ^{11,14}	Once or twice daily	1) CBC – with differential and platelet count – baseline then one (1) to two (2) weeks after each dosage increase, and as clinically indicated 2) Hepatic function panel – baseline and as clinically indicated 3) Pregnancy Test – baseline and as clinically indicated 4) Valproic acid level – 1-2 weeks after initiation and dosage change, then as clinically indicated 5) Serum creatinine and BUN at baseline and as clinically indicated Therapeutic Serum Concentration ¹³ : 50-150 mcg/ml Frequency of Serum Level Monitoring: <ul style="list-style-type: none"> • Day 7-14 • At dosage change • As clinically indicated 	<ul style="list-style-type: none"> • Alopecia • Ataxia • Cognitive impairment • Dizziness • GI upset • Hepatitis • Pancreatitis • Polycystic ovarian syndrome • Rash • Somnolence • Thrombocytopenia • Tremor • Weight gain 	<ul style="list-style-type: none"> • Antipsychotics • Benzodiazepines • Carbamazepine • Lamotrigine¹⁴ • Lithium • MAOIs • Phenytoin • Tricyclic antidepressants • Warfarin

¹¹ Maximum daily dosage should be based upon the medication serum level in the individual patient in the context of clinical response and tolerability.

¹² Recommended dose titration of lamotrigine for patients taking carbamazepine (or other enzyme-inducing drugs) and not taking divalproex: 50mg daily for weeks 1 & 2; 100 mg daily (in divided doses) for weeks 3 & 4; 200 mg daily (in divided doses) for week 5; 300 mg daily (in divided doses) for week 6; up to 400 mg daily (in divided doses) for week 7 and thereafter.

¹³ Therapeutic serum level monitoring of mood stabilizers should be drawn 12-hours after the last dose.

¹⁴ Recommended dose titration of lamotrigine for patients taking divalproex or other forms of valproic acid: 25 mg every other day for weeks 1 & 2; 25 mg daily for weeks 3 & 4; 50 mg daily for week 5; 100mg daily for week 6 and thereafter.

Appendix C: Medication Charts

Adjunctive Agents, Aggression/Hostility

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Lithium <i>Generic available</i> Eskalith® Eskalith® CR	900 mg/day	Check serum level at 3-4 days and adjust (linear kinetics)	900-2400 mg/day	3600 mg/day ¹⁵	Once or twice daily	1) EKG (mandatory for everyone – baseline, yearly and as clinically indicated) 2) CBC – baseline, yearly and as clinically indicated 3) Thyroid studies – baseline; then TSH every 6 months and as clinically indicated 4) BUN, creatinine, glucose and electrolytes; baseline and as clinically indicated 5) UA – baseline and as clinically indicated 6) Pregnancy test – as clinically indicated 7) Lithium Levels – one week after initiation or dosage change and as clinically indicated Therapeutic Serum Concentration ¹⁶ : 0.6-1.5 mEq/L Frequency of Serum Level Monitoring: <ul style="list-style-type: none"> • Day 7 • At dosage change • As clinically indicated 	<ul style="list-style-type: none"> • Acne • Acute renal dysfunction • Cognition • Diarrhea • Dizziness • ECG changes • GI upset • Hypothyroidism • Nausea • Polyuria • Sedation • Thirst • Tremor • Weight gain 	<ul style="list-style-type: none"> • ACE-inhibitors • Caffeine • NSAIDs • Osmotic diuretics • Theophylline • Thiazide diuretics
Oxcarbazepine Trileptal®	600 mg/day	600 mg/day every 7 days	600-2100 mg/day	2400 mg/day	2 - 3 Times Daily	1) Electrolytes – baseline and as clinically indicated 2) Pregnancy test – as clinically indicated	<ul style="list-style-type: none"> • Ataxia • Diplopia • Dizziness • GI upset • Hyponatremia • Somnolence • Tremor 	<ul style="list-style-type: none"> • Antipsychotics • Dihydropyridine calcium antagonists • Oral contraceptive pill • Vitamin D

¹⁵ Maximum daily dosage should be based upon the medication serum level in the individual patient in the context of clinical response and tolerability.

¹⁶ Therapeutic serum level monitoring of mood stabilizers should be drawn 12-hours after the last dose. Therapeutic blood level recommendations based upon data for acute mania in bipolar disorder.

Appendix C: Medication Charts

Adjunctive Agents, Depression

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Citalopram <i>Generic available</i> Celexa®	20 mg/day	10 mg every 2 weeks	20 mg/day	60 mg/day	Once daily in the morning	1) Pregnancy test – as clinically indicated	<ul style="list-style-type: none"> • Agitation • Constipation • Diarrhea • Dizziness • Dry mouth • Fatigue • Headache • Insomnia • Loss of appetite • Nausea • Nervousness • Sexual dysfunction • Somnolence • Sweating 	<ul style="list-style-type: none"> • Clozapine • Cyclosporine • Linezolid • MAOIs • NSAIDs • Pimozide • St. John's Wort • Sympathomimetics • Tramadol • Triptans
Escitalopram Lexapro®	10 mg/day	10 mg every 2 weeks	10-20 mg/day	20 mg/day	Once daily			<ul style="list-style-type: none"> • Cyclosporine • Linezolid • MAOIs • NSAIDs • St. John's Wort • Sympathomimetics • Tramadol • Triptans
Fluoxetine <i>Generic available</i> Prozac®	20 mg/day	10-20 mg every 4 weeks	20 mg/day	40-80 mg/day	Once daily in the morning			<ul style="list-style-type: none"> • Carbamazepine • Clozapine • Cyclosporine • Hydantoins • Linezolid • MAOIs • NSAIDs • St. John's Wort • Sympathomimetics • Thioridazine • Tramadol • Triptans • Tricyclic antidepressants

Appendix C: Medication Charts

Adjunctive Agents, Depression

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Fluvoxamine <i>Generic available</i> Luvox®	50 mg/day	50-100 mg every 2 weeks	100-200 mg/day	300 mg/day	Once or twice daily	See Previous Page	See Previous Page	<ul style="list-style-type: none"> • Carbamazepine • Clozapine • Cyclosporine • Grapefruit • Hydantoins • Linezolid • MAOIs • Methadone • NSAIDs • Ropivacaine • St. John's Wort • Sympathomimetics • Tacrine • Theophyllines • Thioridazine • Tizanidine • Tramadol • Triptans • Tricyclic antidepressants
Paroxetine <i>Generic available</i> Paxil® Paxil CR®	20 mg/day	10-20 mg every 2 weeks	20-30 mg/day	40-60 mg/day	Once daily in the morning			<ul style="list-style-type: none"> • Cyclosporine • Linezolid • MAOIs • NSAIDs • Phenothiazines • St. John's Wort • Sympathomimetics • Tramadol • Triptans • Tricyclic antidepressants
Sertraline <i>Generic available</i> Zoloft®	50 mg/day	50-100 mg every 2 weeks	50-100 mg/day	150-200 mg/day	Once daily in the morning			<ul style="list-style-type: none"> • Carbamazepine • Clozapine • Cyclosporine • Grapefruit • Hydantoins • Linezolid • MAOIs • NSAIDs • Phenothiazines • Pimozide • St. John's Wort • Sympathomimetics • Tramadol • Triptans • Tricyclic antidepressants

Appendix C: Medication Charts

Adjunctive Agents, Depression

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Bupropion <i>Generic available</i> Wellbutrin SR®	150 mg/day	150 mg/day at 3-7 days	225-300 mg/day 200-300 mg/day (SR)	450 mg/day 400 mg/day (SR)	2-3 times daily ≤150 mg/dose Twice daily ≤ (SR)200 mg/dose	1) Pregnancy test – as clinically indicated	<ul style="list-style-type: none"> • Constipation • Dry Mouth • Headache • Insomnia • Nausea • Seizures 	<ul style="list-style-type: none"> • Carbamazepine • Cyclosporine • Linezolid • MAOIs • Ritonavir • Tricyclic antidepressants
Venlafaxine <i>Generic available</i> Effexor® Effexor XR®	37.5 mg/day	37.5 – 75 mg/day every 5-7 days	150-225 mg/day 75-225 mg/day (XR)	375 mg/day	Twice daily Once daily (XR)	1) Pregnancy test – as clinically indicated 2) Blood pressure during dosage titration and as clinically necessary	<ul style="list-style-type: none"> • Anxiety • Blood pressure • Decreased appetite • Dizziness • Dry Mouth • Fatigue • Insomnia • Nausea • Somnolence • Sweating 	<ul style="list-style-type: none"> • Linezolid • MAOIs • St. John's Wort • Sympathomimetics • Tramadol • Triptans
Duloxetine Cymbalta®	30-60 mg/day	None	60 mg/day	60 mg/day	Once or twice daily	1) Pregnancy test – as clinically indicated 2) Blood pressure prior to initiating treatment, during dosage titration, and as clinically indicated 3) Monitor for emergence of suicidal ideation or behavior 4) Hepatic function testing – baseline and as clinically indicated	<ul style="list-style-type: none"> • Anxiety • Blood pressure • Decreased appetite • Dizziness • Dry Mouth • Fatigue • Insomnia • Nausea • Somnolence • Sweating 	<ul style="list-style-type: none"> • Alcohol • Linezolid • MAOIs • St. John's Wort • Sympathomimetics • Thioridazine • Tramadol • Triptans
Mirtazapine <i>Generic available</i> Remeron®	15 mg/day	15 mg every 1-2 weeks	30 mg/day	60 mg/day	Once daily at bedtime	1) Pregnancy test – as clinically indicated	<ul style="list-style-type: none"> • Constipation • Dizziness • Drowsiness • Increased Appetite • Weight Gain • Xerostomia 	<ul style="list-style-type: none"> • Tricyclic antidepressants • Tetracyclic Antidepressants • MAOIs • Sibutramine • Cisapride

Appendix C: Medication Charts

Adjunctive Agents, Depression

Drug	Starting Dose	Titration	Target Dose or Range	Maximum Daily Dose	Schedule	Patient Monitoring Parameters	Side Effects	Selected Drug Interactions
Phenelzine Nardil®	45 mg/day	15 mg/week	60-90 mg/day	90 mg/day	2 - 3 Times Daily	1) Blood chemistries with emphasis on hepatic and renal functions; baseline, yearly and as clinically indicated during prolonged or high dose therapy 2) Pregnancy test – as clinically indicated 3) Blood pressure at baseline and during dosage adjustments and as clinically indicated. Therapeutic ranges for the lab used should be listed on the report	<ul style="list-style-type: none"> • Edema • Insomnia • Orthostatic Hypotension • Sexual Dysfunction • Weight Gain 	<ul style="list-style-type: none"> • Atomoxetine • Bupropion • Carbamazepine • Dextromethorphan • Insulins • Levodopa • Linezolid • Meperidine • SSRIs • St. John's Wort • Sulfonylureas • Sympathomimetics • Tramadol • Triptans • Tricyclic antidepressants • Tyramine foods • Venlafaxine
Tranlycypromine Generic available Parnate®	20-30 mg/day	10 mg/week	20-40 mg/day	60 mg/day	2 - 3 Times Daily			
Amitriptyline Generic available Elavil®	25-75 mg/day	25-50 mg/day every 1-2 days	150 mg/day	300 mg/day	Once daily	1) EKG – baseline and as clinically indicated 2) Pregnancy test – as clinically indicated 3) Blood levels as clinically indicated. • Amitriptyline + Nortriptyline: 120-250 ng/mL ¹⁷	<ul style="list-style-type: none"> • Blurred vision • Constipation • Dry mouth • Orthostatic hypotension • Sedation • Tachycardia • Urinary retention • Weight gain 	<ul style="list-style-type: none"> • Carbamazepine • Ciimetidine • Clonidine • Fluoxetine • Guanethidine • Linezolid • MAOIs • Paroxetine • Procainamide • Quinidine • Quinolones • Rifabutin • Rifampin • St. John's Wort • Sympathomimetics • Valproate • Ziprasidone
Desipramine Generic available Norpramin® Pertofrane®	25-75 mg/day	25-50 mg/day every 1-2 days	150 mg/day	300 mg/day	Once daily	1) EKG – baseline and as clinically indicated 2) Pregnancy test – as clinically indicated 3) Blood levels as clinically indicated. • Desipramine: 100-300 ng/mL ¹⁷		
Doxepin Generic available Sinequan®	25-75 mg/day	25-50 mg/day every 1-2 days	150 mg/day	300 mg/day	Once or Twice daily	1) EKG – baseline and as clinically indicated 2) Pregnancy test – as clinically indicated 3) Blood levels as clinically indicated. • Doxepin + Nordoxepin: 150-250 ng/mL ¹⁷		
Nortriptyline Generic available Pamelor® Aventyl®	25 mg/day	25 mg/day every 2 days	75 mg/day	200 mg/day	Once daily	1) EKG – baseline and as clinically indicated 2) Pregnancy test – as clinically indicated 3) Blood levels as clinically indicated. • Nortriptyline: 50-150 ng/mL ¹⁷		

¹⁷Therapeutic drug monitoring of tricyclic antidepressants can be performed after 5-7 days of consistent dosing. Dose adjustments made to achieve 12-hour blood levels within a therapeutic range.

ADDITIONAL REFERENCES FOR DRUG INFORMATION

- *AHFS Drug Information 2006*, American Society of Health-System Pharmacists, Bethesda, MD, 2006.
- Drug Interactions – <http://medicine.iupui.edu/flockhart/>
- Drug Interactions Facts – <http://online.factsandcomparisons.com/> (subscription required)
- Drug Product Labeling – see specific FDA approved drug prescribing information
- Ketter TA, ed. *Advances in Treatment of Bipolar Disorder*, edited by Terence A Ketter (Review of Psychiatry Series. Volume 24, Number 3; Oldham JM and Riba MB, services editors). Washington, DC: American Psychiatric Publishing. 2005.
- Perry PJ, Alexander B, Liskow BI, Devane CL. *Psychotropic Drug Handbook*, 8th ed. Baltimore, Md: Lippincott Williams and Wilkins, 2007.
- Schatzberg AF, Nemeroff CB, eds. *Textbook of Psychopharmacology*. 3rd ed. Washington DC, American Psychiatric Press Inc. 2003
- Lacy CF, Armstrong LL, Goldman MP, Lance LL. *Drug Information Handbook*, 15th ed. Hudson, OH: Lexi-Comp, 2007.

Appendix D: Side Effect and Co-Existing Symptom Management

Management of Associated or Co-Existing Symptoms in Schizophrenia

Benzodiazepines and other medications with abuse/dependence potential are best avoided in patients with prior history of substance abuse/dependence or who are at risk for substance abuse. Nonaddicting agents are preferred.

<i>Co-Existing Symptom</i>	<i>Recommendations</i>
Agitation/ Excitement	<ul style="list-style-type: none"> - Consider adjunctive medications, including as needed use of oral and intramuscular medications including benzodiazepines, typical antipsychotics, and atypical antipsychotics: <ul style="list-style-type: none"> • Lorazepam 1-4 mg or clonazepam 0.5-2 mg may be used in treating acute agitation. In emergent situations where rapid reduction of agitation is necessary, lorazepam 1-2 mg given intramuscularly may be preferable to oral dosing. The dose may be repeated every 1-2 hours as needed, and onset of effect is generally seen within 15-30 minutes. • Haloperidol 5 mg orally or intramuscularly may be given every 30-60 minutes until patient is calm. • Atypical antipsychotics in intramuscular or oral formulations may be given on an as needed basis to control acute agitation. If oral dosing is used, doses should be initiated at the low end of the dosing range. Intramuscular olanzapine, risperidone oral solution, and intramuscular ziprasidone act more rapidly than their oral counterparts and their use may be warranted in cases where the patient cannot tolerate or does not respond to typical antipsychotic agents and/or benzodiazepines. <ul style="list-style-type: none"> ○ Intramuscular olanzapine 2.5-10 mg, may repeat 2 hours after initial dose and 4 hours after second dose, with a maximum of 30 mg daily. ○ Intramuscular ziprasidone 10-20 mg as needed to a maximum dose of 40 mg daily. The 10 mg dose may be given every 2 hours, and the 20 mg dose may be given every 4 hours. ○ Intramuscular aripiprazole 5.25 – 9.75 mg as needed every two hours to a maximum of 30 mg daily. • Risperidone oral solution is available in 1mg/mL. - Failure of the first trial of pharmacotherapy should be followed by a second trial of an alternative agent above. - After failure of multiple trials of agents to control acute agitation/excitement, consider moving treatment to the next algorithm stage.
Persistent symptoms of Aggression/ Hostility/ Mood Lability	<ul style="list-style-type: none"> - Lithium, valproate and carbamazepine are all therapeutic options for the management of aggression and hostility associated with acute exacerbations in schizophrenia. - If there is no discernible change in the clinical picture after 1-3 weeks, the clinician should discontinue the adjuvant mood stabilizer and consider switching the patient to clozapine.
Depression	<ul style="list-style-type: none"> - Medication treatments for depression in schizophrenia are the same as those used in major depressive disorder. - SSRIs, venlafaxine XR, bupropion SR/XL, duloxetine and mirtazapine are recommended as first line treatments.

<i>Co-Existing Symptom</i>	<i>Recommendations</i>
Insomnia	<ul style="list-style-type: none"> - Promote good sleep hygiene: <ul style="list-style-type: none"> • Encourage regular aerobic exercise at least four hours before bedtime. • Avoid alcoholic beverages. • Encourage regular sleep cycles. • Eliminate noises and distracting lights. • Engage in relaxing activities before bed (reading, sex, meditation, etc.). • Try a glass of milk - Adjunctive medications: <ul style="list-style-type: none"> • Zolpidem 5-10 mg once daily at bedtime. • Zaleplon 5-20 mg (10 mg recommended dose) once daily at bedtime. • Eszopiclone 2-3 mg once daily at bedtime. • Benzodiazepine, such as temazepam 15-30 mg once daily at bedtime or lorazepam 0.5-2mg once daily at bedtime. • Trazodone 25-100mg once daily at bedtime. • Low-dose tricyclic antidepressant, such as amitriptyline 10-50 mg once daily at bedtime. - Brief, targeted cognitive therapy.

Management of Treatment-Emergent Side Effects in Schizophrenia

In general, treatment emergent side effects should be addressed first by dose reduction or medication switching. Prescribing medications for side effects may lead to new side effects. Benzodiazepines and other medications with abuse/dependence potential are best avoided in patients with prior history of substance abuse/dependence or who are at risk for substance abuse. Nonaddicting agents are preferred.

<i>Side Effect</i>	<i>Recommendations</i>
GI Upset	<ul style="list-style-type: none"> - Nausea and diarrhea are usually transient side effects with antidepressants, and improvement should occur within 2-3 weeks after initiation or dose increases. - Administer medication with food and large quantities of liquid. - Consider lowering dose, if possible, or slowing the dose titration. - Persistent GI upset may require changing to an alternative medication or adding an adjunctive agent, such as an H₂ blocker (e.g., famotidine, ranitidine).
Tremor	<ul style="list-style-type: none"> - Enhanced physiologic tremor – A fine tremor of approximately 8-10 Hz; made worse with outstretched hands. <ul style="list-style-type: none"> • Check blood levels of medication, if applicable. • Decrease dose, divide dose, or change to slow release preparation of the medication. • Propranolol can be given at 20-30 mg three times a day. - Parkinsonian tremor – Coarse tremor at rest of approximately 4-6 Hz. <ul style="list-style-type: none"> • See treatment recommendations under Extrapyrarnidal Symptoms (EPS) below.
Sedation	<ul style="list-style-type: none"> - A thorough evaluation of sleep behaviors should be performed, including a patient assessment of sleep quality. - May try dosing medication at bedtime. - Decrease dose if possible. - Substitute a less sedating alternative medication. - Adjunctive medications may be considered. However, in patients with psychosis, adjunctive treatment is not recommended as it may possibly worsen the course of the episode.
Extrapyrarnidal Symptoms (EPS) – Parkinsonian Tremor, Akathisia, and Dystonia	<ul style="list-style-type: none"> - Usually seen with typical antipsychotics or higher doses of risperidone. - Parkinsonian tremor – coarse tremor at rest of approximately 4-6 Hz. <ul style="list-style-type: none"> • Decrease dose, divide dosing, use bedtime dosing, or switch to alternate medication. • Pharmacological treatments include benztropine 1-2 mg twice daily, diphenhydramine 25-50 mg two or three times daily, or propranolol 20-30 mg three times daily. - Akathisia may respond to propranolol 20-30 mg three times a day. If this is not effective, alternatives include clonidine 0.1 mg three times a day, lorazepam 1 mg two or three times a day, or clonazepam 0.5-1mg twice a day. - Dystonic reactions can often be prevented by benztropine 1 mg two or three times a day for the first few days of antipsychotic therapy. Acute dystonic reactions are generally managed with benztropine 1-2 mg IM or lorazepam 1-2 mg IM.

<i>Side Effect</i>	<i>Recommendations</i>
Tardive Dyskinesia	<ul style="list-style-type: none"> - Prescribe typical antipsychotics in the lowest dose necessary for the shortest time possible. Mid-potency typical agents may be preferred if typical antipsychotic is selected. - Use atypical antipsychotic medications. - Consider clozapine which has an extremely low risk of TD. - Consider other treatment modalities, including ECT.
Neuroleptic Malignant Syndrome (NMS)	<ul style="list-style-type: none"> - Patients with a history of NMS should be educated about the need to stay well hydrated and avoid strenuous physical activity when outside during hot weather. - If the patient has been on a FGA, changing to a SGA is reasonable.
Sexual Dysfunction	<ul style="list-style-type: none"> - May consider switching to an alternative medication with lower propensity to cause sexual dysfunction. - If SSRI-induced sexual dysfunction, may consider adding bupropion 75-150mg daily. - Alternatives for the management of sexual dysfunction secondary to psychotropic medications is to add a selective phosphodiesterase (PDE) type 5 inhibitor. Use is contradicted with concurrent nitrates. Caution use with concomitant CYP3A4 inhibitors. Data available for use in females is limited to small, open label trials. <ul style="list-style-type: none"> • Sildenafil 25-100mg one-half to 1 hour before sexual activity. • Tadalafil 10-20mg one-half to 1 hour before sexual activity. • Vardenafil 5-20mg one-half to 1 hour before sexual activity. - Other alternative is cyproheptadine 4-8 mg, given shortly before sexual intercourse. However, cyproheptadine is also a serotonin receptor antagonist, and frequent use in patients with affective symptoms should proceed with caution.

<i>Side Effect</i>	<i>Recommendations</i>
Insomnia	<ul style="list-style-type: none"> - Promote good sleep hygiene: <ul style="list-style-type: none"> • Encourage regular aerobic exercise at least four hours before bedtime. • Avoid alcoholic beverages. • Encourage regular sleep cycles. • Eliminate noises and distracting lights. • Engage in relaxing activities before bed (reading, sex, meditation, etc). • Try a glass of warm milk. - If due to concomitant antidepressant use: reduce the dose of antidepressant, if possible. - Try moving the dosing of the medication to the morning. - Adjunctive medications: <ul style="list-style-type: none"> • Zolpidem 5-10 mg once daily at bedtime. • Zaleplon 5-20 mg (10 mg recommended dose) once daily at bedtime. • Eszopiclone 2-3 mg once daily at bedtime. • Benzodiazepine, such as temazepam 15-30 mg once daily at bedtime or lorazepam 0.5-2mg once daily at bedtime. • Trazodone 25-100mg once daily at bedtime. • Low-dose tricyclic antidepressant, such as amitriptyline 10-50 mg once daily at bedtime. - Brief, targeted cognitive therapy.
Weight Gain	<ul style="list-style-type: none"> - Exercise (walking, jogging, swimming) for at least 3 times weekly, and for at least 30 minutes each time. - Diet: <ul style="list-style-type: none"> • Eat smaller portions of 3 meals per day. • Decrease excess fats (decrease fried foods, eat lean meats, increase vegetables, salads, and fruits). • Decrease excessive low nutritional content carbohydrate (soft drinks, desserts, candy, gravies, potatoes, white bread). - Avoid snacking, and particularly no evening snacks.

Appendix E: TMAP Publications

- Altshuler KZ, Rush AJ. Computerizing the Texas Medication Algorithm Project. *Behavioral Healthcare Tomorrow*. 1999;60:284-91.
- Chiles JA, Miller AL, Crismon ML, Rush AJ, Krasnoff AS, Shon S. The Texas Medication Algorithm Project. Development and implementation of the schizophrenia algorithm. *Psychiatric Services* 1999;50:69-74.
- Chuang WC, Crismon ML. Evaluation of a schizophrenia medication algorithm in a state hospital. *American Journal of Health-System Pharmacy* 2003;60:1459-67.
- Crismon ML, Buckley PF. Schizophrenia. In: DiPiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM, eds. *Pharmacotherapy: a pathophysiologic approach, 6th ed*. New York, NY, McGraw Hill, 2005:1209-1233.
- Crismon ML, Dorson PG. Schizophrenia. In: DiPiro JT, Talbert RL, Yee GC, Matzke GR, Wells BG, Posey LM, eds. *Pharmacotherapy: a pathophysiologic approach, 5th ed*. New York, NY, McGraw Hill, 2002;1219-1242.
- Dewan NA, Conley D, Svendsen D, Shon SP, Staup JR, Miller AL, Crismon ML, Rush AJ, Trivedi M, Skale T, Keck PE, Strawkowski SM. A quality improvement process for implementing the Texas Algorithm for schizophrenia in Ohio. *Psychiatric Services* 2003;54:1646-49.
- Gilbert DA, Altshuler KZ, Rago WV, Shon SP, Crismon ML, Toprac MG, Rush AJ. Texas Medication Algorithm Project: Definitions, rationale and methods to develop medication algorithms. *Journal of Clinical Psychiatry* 1998;59:345-51.
- Kashner TK, Rush AJ, Crismon ML, Toprac M, Carmody TJ, Miller AL, Trivedi M, Wicker A, Suppes T. An empirical analysis of cost-outcomes of the Texas medication algorithm project. *Psychiatric Services* 2006;57:648-659.
- Kashner TM, Carmody TJ, Suppes T, Rush AJ, Crismon ML, Miller AL, Toprac M, Trivedi M. Catching-up on health outcomes: The Texas Medication Algorithm Project. *Health Services Research* 2003;38:311-31.
- Kashner TM, Rush AJ, Surís A, Biggs MM.; Gajewski VL, Hooker DJ, Shoaf T, Altshuler KZ. Impact of Structured Clinical Interviews on Physician Behavior in Community Mental Health Settings. *Psychiatric Services* 2003;54:712-718.
- Mellman TA, Miller AL, Weissman E, Crismon ML, Essock SM, Marder SR. Evidence based medication treatment for severe mental illness: a focus on guidelines and algorithms. *Psychiatric Services* 2001;52:619-625.
- Miller AL, Chiles JA, Chiles J, Crismon ML. *TIMA procedural manual: schizophrenia algorithm*. Austin, TX: Texas Department of Mental Health and Mental Retardation, 2000. 59 pp.
- Miller AL, Chiles JA, Chiles J, Crismon ML. *TMAP procedural manual: Schizophrenia module physician manual*. Austin, TX: Texas Department of Mental Health and Mental Retardation, 1998. 59 pp.

- Miller AL, Chiles JA, Chiles JK, Crismon ML, Rush AJ, Shon SP. The TMAP schizophrenia algorithms. *Journal of Clinical Psychiatry* 1999;60:649-657.
- Miller AL, Crismon ML, Hall CS, Baker CB. Advances in psychopharmacology and pharmacoconomics. In: Levin B, Petrila J, (eds.) *Mental health services: a public health perspective, 2nd ed.* New York: Oxford University Press, 2004:344-70.
- Miller AL, Crismon ML, Rush AJ, Chiles J, Kashner M, Toprac M, Carmody T, Biggs M, Shores-Wilson K, Chiles J, Witte B, Bow-Thomas C, Velligan DI, Trivedi M, Suppes T, Shon S. The Texas Medication Algorithm Project: clinical results for schizophrenia. *Schizophrenia Bulletin* 2004;30:627-47.
- Miller AL, Dassori A, Ereshefsky L, Crismon ML. Recent issues and developments in antipsychotic use. In: Dunner DL, Rosenbaum JF, eds. *The Psychiatric Clinics of North America Annual Review of Drug Therapy 2001.* Philadelphia, PA: W.B. Saunders Company, 2001;8:209-235.
- Miller AL, Hall CS, Buchanan RW, Buckley PF, Chiles J, Conley RR, Crismon ML, Ereshefsky L, Essock SM, Finnerty M, Marder SR, Miller D, McEvoy J, Rush AJ, Saeed SA, Schooler NR, Shon S, Stroup S, Tarin-Godoy B. The Texas Medication Algorithm Project antipsychotic algorithm for schizophrenia: 2003 Update. *Journal of Clinical Psychiatry* 2004;65:500-8.
- Miller AL, Hall CS, Crismon ML, Chiles J,. *TIMA procedural manual: schizophrenia algorithm.* Austin, TX: Texas Department of Mental Health and Mental Retardation, 2003. 77 pp.
- Miller, A, Craig, CS. Algorithms for the treatment of schizophrenia: An absolute need. *The Journal of Psychotic Disorders: Reviews & Commentaries*, 5(3):3, 2001.
- Moore TA, Buchanan RW, Buckley PF, Chiles JA, Conley RR, Crismon ML, Essock SM, Finnerty M, Marder SR, Miller DD, McEvoy JP, Robinson D, Schooler NR, Shon SP, Stroup TS, Miller AL. The Texas Medication Algorithm Project Antipsychotic Algorithm for Schizophrenia: 2006 Update. *J Clin Psychiatry.* 2007; 68:1751-1762.
- Parkman CA, Evans KJ, Cox R. Mapping out care with TMAP. *Nursing* 2000;30:74.
- Rago WV, Shon SP. "The Texas Medication Algorithm Project." In *Improving Mental Health Care. Commitment to Quality*, Dickey B and Sederer LI (eds). Washington, DC: American Psychiatric Publishing, Inc., 2001.
- Rush AJ, Crismon ML, Kashner TM, Toprac MG, Carmody TJ, Trivedi MH, Suppes T, Miller AL, Biggs MM, Shores-Wilson K, Witte B, Shon SP, Rago WV, Altshuler KZ. Texas Medication Algorithm Project – Phase III (TMAP-III): rationale and design. *Journal of Clinical Psychiatry* 2003;64:357-69.
- Rush AJ, Crismon ML, Toprac MG, Shon SP, Rago WV, Miller AL, Suppes T, Trivedi MH, Biggs MM, Shores-Wilson K, Kashner TM, Altshuler KZ. Implementing guidelines and systems of care: Experiences with the Texas Medication Algorithm Project (TMAP). *Journal of Practical Psychiatry & Behavioral Health* 1999;5:75-86.
- Rush AJ, Rago B, Toprac M, Crismon ML, Stone S, Shon S. Toward the development of medication algorithms for the treatment of major depressive, schizophrenic, and bipolar disorders - a white paper. Texas Department of Mental Health and Mental Retardation, Austin, TX, 1996, 29 pp.

- Rush AJ, Rago WV, Crismon ML, Toprac MG, Shon SP, Suppes T, Miller AL, Trivedi MH, Swann A, Biggs MM, Shores-Wilson K, Kashner M, Pigott TA, Chiles CA, Gilbert DA, Altshuler KZ. Medication treatment for the severely and persistently mentally ill: The Texas Medication Algorithm Project. *Journal of Clinical Psychiatry* 1999;60:284-291.
- Shon SP, Crismon ML, Toprac MG, Trivedi M, Miller AL, Suppes T, Rush AJ. Mental health care from the public perspective: The Texas Medication Algorithm Project. *Journal of Clinical Psychiatry* 1999;60 (suppl 3):16-20.
- Shon SP, Toprac MG, Crismon ML, Rush AJ. Strategies for implementing psychiatric medication algorithms in the public sector. *Journal of Practical Psychiatry & Behavioral Health* 1999;5:32-36.
- Shores-Wilson K, Biggs MM, Miller AL, Carmody TJ, Chiles JA, Rush AJ, Crismon ML, Toprac MG, Witte BP, Webster JC. Itemized clinician ratings versus global ratings of symptom severity in patients with schizophrenia. *International Journal of Methods in Psychiatric Research* 2002;11:45-53.
- Starkweather K, Shon SP, Crismon ML. A Texas-sized prescription: providers report progress with medication guidelines. *Behavioral Healthcare Tomorrow* 2000;9(4):44-46.
- Starkweather K, Shon SP. Implementing Medication Algorithms: The Texas Experience. *Psychiatrist Administrator* 2001;1:13-18.
- Toprac MG, Dennehy EB, Carmody TJ, et al. Implementation of the Texas Medication Algorithm Project Patient and Family Education Program. *J Clin Psychiatry*. 2006;67:1362-1372.
- Toprac MG, Rush AJ, Conner TM, Crismon ML, Dees M, Hopkins C, Rowe V, Shon SP. The Texas Medication Algorithm Project patient and family education program: a consumer-guided initiative. *Journal of Clinical Psychiatry* 2000;61:477-486.
- Trivedi MH. Using treatment algorithms to bring patients to remission. *Journal of Clinical Psychiatry* 2003;64 (Suppl 2):8-13.
- Trivedi, MH, Kern, J.K., Baker, S.M. and Altshuler, K.Z. Computerizing medication algorithms and decision support systems for major psychiatric disorders. *Journal of Psychiatric Practice*, 2000; 6:237-246.
- Velligan DI, DiCocco M, Bow-Thomas C, Cadle C, Glahn DC, Miller AL, Biggs MM, Shores-Wilson K, McKenzie CA, Crismon ML. A brief cognitive assessment (BCA) for use with schizophrenia patients in a community clinic. *Schizophrenia Research* 2004;71:273-83.

Appendix F: Minimum Data Set for Documentation

The following information should be entered on the Clinical Record Form at each patient visit:

1. Patient identification information

Indicate information required by the health care organization.

2. Date

Date of visit (month/day/year)

3. Service activity code

Service activity or billing code for this visit

4. Physician/clinician code or identification

5. Duration of visit

Record start and end times of visit (hour:minute am/pm).

6. Current diagnoses

Record the current psychiatric diagnoses using DSM IV-TR codes. Please place primary diagnosis first.

7. Current algorithm

Check box of the specific algorithm that is being used.

8. Current stage in algorithm and weeks in this current stage

Record current stage in algorithm at the beginning of this visit and how many weeks the patient has been in this stage.

9. Vital signs

Record current vital signs: weight, height, blood pressure, pulse rate.

10. Most recent drug levels

Most recent values (as applicable) with date

11. Has patient taken medications as prescribed?

Check appropriate box.

12. Any other medications taken during the past week?

Include any prescriptions, over-the-counter medications, or complementary medications taken in addition to medications prescribed by this physician.

13. Patient global self report

Record patient's results, including symptom severity and side effects.

14. Clinical rating scales

Record the scores of any and all appropriate clinical rating scales, including POS SX, NEG SX, QIDS (SR or C), BDSS, AIMS, and any others. Although only the total score is required for the Minimum Data Set, greater clinical utility is achieved by listing each item score for the scale or scales used. The individual rating scale items can be preprinted on the CRF if desired.

These items provide a global impression of the clinician's impression of the severity of each of these symptoms as observed at the visit as well as during the week prior to the visit.

For items 15 – 17, a scale of 0 – 10 should be used:

0 = No symptoms

5 = Moderate symptoms

10 = Extreme symptoms

15. Core symptoms

These are the severity of the core symptoms for the three adult disorders for which algorithms have been developed: mania, depression, positive psychotic symptoms, and negative symptoms.

16. Other symptoms

These include other symptoms that are commonly seen in individuals with mental disorders and include: irritability, mood lability, agitation, anxiety, level of interest, appetite, energy, and insomnia. A space is left in case the clinician wishes to add additional symptoms that may be present in a given patient.

17. Overall side effect severity

Rate the overall level of side effect severity from all medications being taken by the patient.

18. Suicidal or homicidal

Indicate if the patient is presently suicidal or homicidal and, if yes, please comment in the progress note section.

19. Overall functioning

Rate from 0 – 10 (0 = Low and 10 = High) your overall impression of the patient's ability to function on a daily basis. Please note: this is not a GAF score, but the clinician's overall impression of how the patient has been functioning during the last week.

20. Are serum concentrations needed?

This provides a prompt for the clinician to order medication serum concentrations if they are needed. If yes, please specify in the progress note section.

21. Rationale for diagnostic and other services

The rationale for ordering diagnostic and other services should be clearly documented.

22. Medication response

Please indicate the patient's response to the medication since the beginning of the current stage. Check the box that applies. Please note that this is medication response and, depending on comorbidity and the patient's psychosocial situation, this may not necessarily represent the patient's overall improvement in mental health status.

23. Rationale for change in medication

If medication is being changed (including dose changes), please note rationale by checking all boxes that apply.

24. Prescription information

- This information should be completed regardless of whether a patient is getting a new prescription for ongoing medications.

- List all medications being taken by the patient for the core syndrome, other symptoms, or side effects.
- Indicate via check mark, if this is a new medication, continuation of a previous medication, or medication being discontinued at this visit.
- Provide the following information: dose, frequency, duration the medication is to be taken, titration (or tapering) schedule, and any other pertinent information describing the medication or use of this medication.
- Indicate via check mark the following:
 - S = Core symptoms
 - OS = Other symptoms
 - SE = Side effects of S or OS medications

25. Progress note

Use the progress note to indicate additional information, assessments, or impressions not addressed elsewhere or to expand on information already given. This section should also address any variation from algorithm-based treatment. Clinics may use preprinted templates for this section if they wish.

26. Next visit

The treating clinician indicates the recommended number of weeks until the patient should return to the clinic. Clinic staff should record the actual date of the next scheduled visit.

27. Signature and title

Treating clinician should sign name and degree designation or title.